


CATALOGUE



AIZ INTERNATIONAL CO., LTD

Veterinary medicine



www.aizinternational.com





AIZ INTERNATIONAL CO., LTD

Quality affirms brand



AIZ INTERNATIONAL CO., LTD

INTRODUCTION

AIZ Company is one of the professional suppliers of veterinary medicine for livestock, poultry, aquatic and health care products in Vietnam.

All our products are manufactured in factory with international standard - GMP (Good Manufacturing Practices), GLP (Good Laboratory Practice), GSP (Good Storage Practices).

With 5 modern production lines, standard laboratory system, research and development department, and a team of highly qualified employees - AIZ is constantly striving to produce good quality products.

AIZ Company is proud of the distribution network covering all provinces and cities nationwide. Not only that, AIZ's products are available in many markets around the world.



With the motto "Quality affirms brand", AIZ constantly develops and enhances product quality to satisfy customers' demand. Besides, AIZ always innovates and promises with many progress in the future.



Quality affirms brand

TABLE OF CONTENTS

No.	Content	Page	No.	Content	Page	No.	Content	Page	No.	Content	Page
I. SUPPORTIVE PREVENTION& TREATMENT, SUPPLYMENT			III. ANTI-INFLAMMATION, PAIN RELIEF, ANTIPIRETTIC			IV. ANTIBIOTIC			V. ANTI-PARASITES, ANTI-COCCIDIOSIS, ANTI-FUNGI		
Injectable solution			Injectable solution			Intramary suspension			Injectable solution		
1.	ABUSAL.....	1	1.	ANAZIN.3000.....	12	1.	MASTI-CURE.....	22	1.	IVERMAX.....	33
2.	ADE MIJECT.....	1	2.	BUTAZONE-Z.....	12	2.	CEFAOCIN.....	22	2.	MECZ.....	33
3.	AMINOPLUS.....	1	3.	CORHENYL.....	12	3.	LEAXAMIN.....	22	3.	NITROVET.....	33
4.	CIUMY.....	1	4.	CORNIUM.....	12	4.	TIOFUR DRY.....	22	4.	SANTEL MEC.....	33
5.	ENERGY PLUS.....	1	5.	CORTYL VET.....	12	5.	TIOFUR LC.....	22	5.	SULONTIN.....	33
6.	GLUCONAMIC KC.....	1	6.	CYNOLON.....	12	Oral solution/ suspension			Oral solution		
7.	HAEVET JECT.....	2	7.	DECOXASONE.....	13	1.	CYLOSIN.....	23	1.	COCCIVET.....	33
8.	HEXIN Z.....	2	8.	DECOZ.....	13	2.	DOHEFLOR.....	23	2.	COCUS TREAT.....	34
9.	IRON B12.....	2	9.	DEROZI.....	13	3.	ENROHEXIN.....	23	3.	COSISTRE.....	34
10.	LEN-E VITA.....	2	10.	DIPYHEXIN.....	13	4.	ENROVET.....	23	4.	DAZONE.....	34
11.	LIVETECT.....	2	11.	DIPYRON+C.....	13	5.	FLOTYL.....	23	5.	DEWOM.....	34
12.	MULBINOL.....	2	12.	FLUCOID.....	13	6.	LOSINVET.....	23	6.	SUPERIL.....	34
13.	PYTO.....	3	13.	FLUMEA.....	14	7.	NEO-S.....	24	7.	WOR-F.....	34
14.	RACING VET.....	3	14.	KEFEN.....	14	8.	PEC-COLI.....	24	External solution		
15.	SORBILEX.....	3	15.	NIXIN-F.....	14	9.	PECTYL.....	24	1.	VEROTIN.....	35
16.	SUQIMIN.....	3	16.	SOLINE-Z.....	14	10.	TREAVET.....	24	2.	XIMO.....	35
17.	TAPHOMIN.....	3	17.	TRYFASIN.....	14	12.	VATIL-ORAL.....	24	Powder		
18.	THIRON.....	3	Oral solution			Powder for injection			1.	ADE SULTRY.....	35
19.	VIT B JECT.....	4	1.	TRYPFA.....	14	1.	STREPTONIN.....	25	2.	COX-STOP.....	35
20.	VITAMIN C.2000.....	4	Powder			2.	TIOFURDIUM.....	25	3.	IVER-WS.....	35
21.	VITAMIX.....	4	1.	DEROZI-PR.....	15	3.	TRIXONE.....	25	4.	LEVAZOL.....	35
Oral solution			2.	METAMIC.....	15	Powder			5.	NYTANZ.....	36
1.	CALCI-PLUS.....	4	3.	PARA+C.....	15	1.	AMOXY-WS.....	25	6.	PROLICOX.....	36
2.	CALYP AZ.....	4	4.	PYRIVET.....	15	2.	AUCOLIN.....	25	7.	PROXALIN.K.....	36
3.	E-SEL LIQUID.....	4	5.	SALIC.....	15	3.	BERINTREP-WS.....	25	8.	VAMISONE.....	36
4.	GALYVET.....	5	IV. ANTIBIOTIC			4.	BIOCILIN CS.....	26	9.	VERYDINE.....	36
5.	HAELTHVET.....	5	Injectable solution			5.	CEPLYTYN.....	26	10.	ZOBENVET.....	36
6.	HERBAVET.....	5	1.	COLISTIN.....	16	6.	COLIMAX-S.....	26	VI. DISINFECTANT		
7.	HEXIN.....	5	2.	COLYXA.....	16	7.	COLIVIT.TETRA.....	26	1.	ANTIBACTERIAL SPRAY.....	37
8.	MEPHO.....	5	3.	COSPIRA.....	16	8.	COLYDOF.....	26	2.	DYTAN.....	37
9.	MINOCID.....	5	4.	DECOGEN.....	16	9.	COLYN-Z.....	26	3.	OXIDAN.....	37
10.	PHOCIUM.....	6	5.	DOF-SONE.....	16	10.	COMPLEX AMIN.....	27	4.	VODINE.....	37
11.	PROLIVER.....	6	6.	DYPATE.....	16	11.	CYCOSIN.....	27			
12.	SOBICAR.....	6	7.	FLOCOVET.....	17	12.	CYN-DOX.....	27			
13.	TOXYLIQ.....	6	8.	FLORJECT.....	17	13.	DOXLOSIN.....	27			
14.	VITAMI COMPLEX.....	6	9.	FLOTYL.....	17	14.	DOXYFLOR.....	27			
Powder			10.	LINCO S.1500.....	17	15.	DOXYVET.....	27			
1.	ACIVITA.....	6	11.	MARCINE.....	17	16.	ERYCIN-B.....	28			
2.	ADE THIAM.....	7	12.	MAXLOX.....	17	17.	GENTACIN.....	28			
3.	ASCOBIC-WS.....	7	13.	MEGNICOL.....	18	18.	LEXYN.....	28			
4.	BLOOVET.....	7	14.	MULIN-CS.....	18	19.	MAXFOC-WS.....	28			
5.	DETOXVET.....	7	15.	OXYTRA-LA.....	18	20.	MOXCYCIN.....	28			
6.	DIGEST-TONIC.....	7	16.	QUINO-MAX.....	18	21.	NEOCIN-WS.....	28			
7.	GINSENG.....	7	17.	SPYCO.....	18	22.	NEOTYN.....	29			
8.	HAEZET.....	8	18.	TETRA FLU.....	18	23.	NEOX.....	29			
9.	HEALBALAN.....	8	19.	THIAM-LOSYN.....	19	24.	NITAXY.....	29			
10.	HEASUPPORT.....	8	20.	THIAM-SONE.....	19	25.	NIXOLIN.....	29			
11.	MILK ENHAN.....	8	21.	TILSYN.....	19	26.	OXIMIN.....	29			
12.	PEROL WS.....	8	22.	TYLANVET.....	19	27.	OXPATE.....	29			
13.	PROBI-ALLY.....	8	23.	TYLO-TETRA.....	19	28.	PECLYN.....	30			
14.	PROBIZ.....	9	24.	TYLO-GENTA.....	19	29.	SAMYCIN.....	30			
15.	PROTIC.....	9	Injectable suspension			30.	SAMYPRIM.....	30			
16.	RACE COMPE.....	9	1.	AMGEN-S.....	20	31.	SASIN.....	30			
17.	REHYDRATE.....	9	2.	AMPI-CS.....	20	32.	SULAM.....	30			
18.	RINEVET.....	9	3.	CEFOXYL.....	20	33.	TERAC.....	30			
19.	SORBIVET.....	9	4.	CEFTIFEN.....	20	34.	TETROZ.....	31			
20.	SUPPLEVET.....	10	5.	CEQUIN 4.....	20	35.	THOXAPRY.....	31			
21.	TAVILEX.....	10	6.	CETIMAX.....	20	36.	TIMIVET.....	31			
22.	TROLYT AIZ.....	10	7.	CLA-MOX.....	21	37.	TRACINVET.....	31			
23.	VIRALMIN.....	10	8.	COLIMOX.....	21	38.	TYCOCIN.....	31			
24.	VITA-CA.....	10	9.	NOVAMOX-LA.....	21	39.	TYLAGEN.....	31			
25.	VITASUPPLY.....	10	10.	PENISTREP.....	21	40.	TYLANVET.....	32			
II. HORMONES			11.	TRIME-ZONE.....	21	41.	TYLOTETRA.....	32			
1.	ETD.....	11	12.	TRIXON.....	21	42.	VIT-NEO.....	32			
2.	ETD VIT.....	11				43.	VOTYL-Z.....	32			
3.	TOXYL.....	11									



Injectable solution

ABUSAL

50 ml, 100 ml



COMPOSITION: 100 ml contains
 1-(n - Butylamino) - 1 methylethyl -
 Phosphonic acid.....15,000 mg
 Cyanocobalamin.....10,000 mcg

INDICATIONS:
 - Enhance metabolism, increasing weight.
 - Prevent stress, high recover after diseases.

DOSAGE :
 - Large ruminant: 1 ml/ 20 - 30 kgs B.W.
 - Poultry: 1 ml/ 4 kgs B.W.

COMPOSITION: 100 ml contains
 Vitamin A.....8,000,000 IU
 Vitamin D₃.....4,000,000 IU
 Vitamin E.....2,000 mg

INDICATIONS:
 Stimulate appetite. Increase fertility.
 Enhance absorption of Ca, P for fetus
 development, prevent poor growth.

DOSAGE:
 I.M administration
 - Piglets, dogs, cats: 1 - 2 ml/ injection.
 - Swine, calves, sheep, goats: 4ml/ injection.
 - Sows, boars: 6 ml/ injection.

ADE MIJECT

50 ml, 100 ml



AMINOPLUS

50 ml, 100 ml



COMPOSITION: 1 ml contains
 Dextrose.....50 mg Pyridoxine HCl...0.1 mg
 Calcium chloride.....2 mg Nicotinamide.....3 mg
 Potassium chloride....2 mg Thiamine HCl.....0.1 mg
 Sodium chloride.....7.5 mg Methionine.....0.525 mg
 Magnesium chloride...2 mg Lysine HCl.....0.525 mg
 Riboflavin.....0.05 mg Tryptophan.....0.715 mg
 D-pantothenol.....0.1 mg Threonine.....0.35 mg
 Excipient (Isoleucine, Phenylalanine, Valine, Leucine)
 q.s.....1 ml

INDICATIONS:
 Resistance aid, anti-stress. Recovery from
 disease, prevention of asthenic. Gain weight
 & reproductivity aid.

DOSAGE:
 I.M, I.V administration
 - Piglets, dogs, cats: 1ml/ 10 kgs B.W.
 - Swine, bovine: 1 ml/ 15 - 25 kgs B.W.
 - Poultry: 1 ml/ 1 liter of water.

COMPOSITION: 100 ml contains
 Calcium gluconate.....10 g
 Vitamin B₁₂.....20,000 mcg

INDICATIONS:
 Treatment of milk fever, convulsions,
 paralysis, poor growth and anemia.

DOSAGE:
 For oral administration or slow I.M, I.V
 injection
 - Cattle: 30 - 40 ml/ head/ day.
 - Poultry: 1 ml/ 6 - 10 kgs B.W.

CIUMY

50 ml, 100 ml



ENERGY PLUS

50 ml, 100 ml



COMPOSITION: 1 ml contains
 ATP.....6.5 mg
 L.arginine.....1 mg
 L.carnitine.....1 mg
 Pangamate.....275 mg

INDICATION:
 Energy supplement, fast health recovery.
DOSAGE:
 For intramuscular injection only
 Fighting roosters: 0.4 ml/ kg B.W.

COMPOSITION: 100 ml contains
 Calcium gluconate.....25,000 mg
 Vitamin C.....500 mg
 Vitamin K₃.....500 mg
 Tolfennamic acid.....3,500 mg

DOSAGE:
 I.M. or S.C. administration for 3 - 5 days
 - Cattle, buffalos: 1 ml/ 12 - 15 kgs B.W.
 - Swine, goats, sheep: 1 ml/ 7 - 10 kgs B.W.
 - Poultry: can be used by oral route: 1 ml/ 5 - 6 kgs B.W.

INDICATIONS:
 - Supplement of glucose, vitamins for
 metabolism, supplement of energy for animal.
 - Rapid health recovery after diseases, giving
 birth or depression
 - Antipyretic, anti - inflammation and
 supportive treatment.

GLUCONAMIC KC

50 ml, 100 ml





HAEVET JECT

50 ml, 100 ml



COMPOSITION: 100 ml contains
 Vitamin K₃.....1 g
INDICATIONS:
 Hemorrhage, enteritis, hepatitis,
 coccidiosis with *E.coli*, injury, trauma.

DOSAGE:
 I.M administration
 - Swine: 1 ml / 10 kgs B.W.
 - Chickens: 1 ml / 5-10 kgs B.W.
 - Buffalos, cattle, sheep, goats: 1 ml / 10 - 20 kgs B.W

COMPOSITION: 100 ml contains
 Bromhexin HCl.....1,000 mg
INDICATIONS:
 Reduce cough, bronchus mucus and
 excrete phlegm.

DOSAGE:
 - Swine, goat, sheep, dogs, cats: 1 ml / 25 -
 30 kgs B.W.
 - Buffalos, cattle: 1 ml / 30 - 35 kgs B.W.

HEXIN Z

50 ml, 100 ml



IRON B12

50 ml, 100 ml



COMPOSITION: 100 ml contains
 Fe (as dextran complex).....10 g
 Vitamin B₁₂.....10 mg
INDICATIONS:
 Prophylactics and treatment of anemia.

DOSAGE:
 For I.M or S.C administration
 - Calves, goats, sheep: 1 ml / 10 kgs B.W
 S.C, in the first week of age (4 - 8 ml / head).
 - Piglets: 2 ml I.M, first 3 days of age.

COMPOSITIONS: 1 ml contains
 Vitamin E.....50 mg
 Sodium selenite.....0.5 mg
INDICATIONS:
 Treatment and prevention of disease
 associated to vitamin E and selenium
 deficiency as encephalomalacia, infertility in
 livestock.

DOSAGE:
 For I.M, S.C administration
 - Calves, goats, sheep: 2ml / 10 kgs B.W.
 - Swine: 1ml / 10 kgs B.W.

LEN E-VITA

50 ml, 100 ml



LIVETECT

50 ml, 100 ml



COMPOSITION: 1 ml contains
 DL-Acetyl-Methionine.....200 mg
INDICATION:
 Treatment of liver diseases.

DOSAGE:
 I.V, I.M, S.C injection or oral administration
 - Cattle: 50-100 ml.
 - Horse: 50 ml
 - Small ruminants: 20 ml
 - Dogs: up to 10 ml

COMPOSITION: 100 ml contains
 Vitamin B₁.....1 g
 Vitamin B₂.....400 mg
 Vitamin B₆.....400 mg
 Vitamin B₁₂.....1,000 mcg
 Nicotinamide.....5,000 mg
 D - panthenol.....500 mg

INDICATIONS:
 Prevention or treatment of vitamin B
 deficiencies, stress.
DOSAGE:
 For S.C or I.M administration
 - Swine: 0.5 - 1 ml / 10 kgs B.W.
 - Cattle, horse: 1 ml / 15 - 20 kgs B.W.

MULBINOL

50 ml, 100 ml



Injectable solution



COMPOSITION: 100 ml contains
 Atropin sulfate.....100 mg
INDICATIONS:
 Treatment of intestinal spasm, severe diarrhea, tracheal-bronchial spasm.

DOSAGE:
 S.C or I.M administration
 - Dogs, cats: 1 ml / 25 - 30 kgs B.W.
 - Swine, sheep, goats: 1 ml / 20 - 25 kgs B.W.
 - Buffalo, bovine: 1 ml / 15 - 20 kgs B.W.

COMPOSITION: 1 ml contain
 Sodium bicarbonate.....8.4%
 U.T.P.....15 mg
 A.T. P.....20 mg
 L.arginine HCl.....22 mg
DOSAGE:
 5 - 10 ml per day for 2 days before the competition, including the day of the race.
 Slow I.V.

INDICATION:
 Preparation for the test. Dystrophies and myalgia. Physical exhaustion. Overtraining. Neuromuscular tonic.



COMPOSITION: 1 ml contains
 Sorbitol.....130 mg
 Methionine.....50 mg
 Lysine.....50 mg
 Vitamin B₁₂.....200 mcg
 Vitamin B₆.....20 mg
 Vitamin B₂.....2 mg

INDICATIONS:
 Increasing bile secretion to stimulate digestion, increasing growth rate.
DOSAGE:
 I.M injection, 1 time/ day for 3 - 5 days
 - Swine, goats: 1 ml / 10 - 20 kgs body weight.
 - Chickens, ducks: 1 ml / 5 - 10 kgs body weight.
 - Buffalos, cattle: 1 ml / 25 - 30 kgs body weight.
 - Prevention, growth: Inject 1 time / 1 - 2 weeks.

COMPOSITION: 1 ml contains
 Lactose.....50 mg
 Vitamin B₁.....10 mg
 Methionin.....50 mg
 Sorbitol.....100 mg
INDICATIONS:
 Stimulation of milk secretion. Anti-depression, anti-constipation. Detoxication, stimulation of digestion.

DOSAGE:
 I.M administration
 Once time per day, 3 - 5 days after parturition
 - Sows: 10 ml / sow / time.
 - Cows: 20 ml / sow / time.



COMPOSITIONS: 100 ml contains
 Butaphosphan.....15,000 mg
 Tauribe.....3,730 mg
 Methionine.....1,870 mg
 Vitamin B₁₂.....7,500 mcg
 Nicotinamide.....2,300 mg
INDICATIONS:
 Treatment of acute and chronic metabolism disorder. Increasing growth rate in young animal Aid resistance, high recovery.

DOSAGE: I.M, I.V or S.C administration.
 - Buffalos, cattle, horse: 5 - 25 ml
 - Swine: 2.5 - 10 ml
 - Piglets: 1 - 2.5 ml
 - Calves, sheep, pony: 5 - 12 ml
 - Sheep: 2.5 - 5 ml
 - Young sheep: 1.5 - 2.5 ml
 - Dogs: 0.5 - 5 ml
 - Cats: 0.5 - 2.5 ml
 - Poultry: Dilute into water: 0.3 - 0.5 ml / individual

COMPOSITION:
 Thiamine hydrochloride.....500 mg / 1.7 cc
INDICATION:
 Prevention & treatment of vitamin B₁ deficiency.

DOSAGE:
 I.V, I.M administration
 - Horses: 1 - 2 ml.
 - Cattle, camel: 3 - 4 ml.
 - Swine, lambs, sheep: 2 - 50 ml / day.
 - Dogs & cats: 2 - 20 ml / day.



VIT B JECT

50 ml, 100 ml



COMPOSITION: 100 ml contains
Cyanocobalamin.....50,000 mcg
INDICATION:
Prevention and treatment of anemia due to lack of Vitamin B₁₂, malnutrition.

DOSAGE:
I.M and S.C injection
- Piglet: 1 ml/ head.
- Pig: 2 - 4 ml/ head.
- Bovine: 5 - 8 ml/ head.
- Calves: 3 - 5 ml/ head.

COMPOSITION: 100ml contains
Acid ascorbic.....20,000 mg
DOSAGE:
1 ml/ 15 - 20 kgs B.W.

INDICATION:
Vitamin C supplement. Influenza treatment and stress prevention. Enhanced resistance, quick recover from disease

VITAMIN C.2000

50 ml, 100 ml



VITAMIX

50 ml, 100 ml



COMPOSITION: 100 ml contains
Vitamin A.....50,000 IU
Vitamin D₃.....10,000 IU
Vitamin E.....10 mg
Vitamin B₁.....2.5 mg
Vitamin B₆.....1.5 mg
Vitamin C.....25 mg
Nicotinamide.....17.5 mg
Vitamin K₃.....1.5 mg

INDICATIONS:
Stimulation appetite, prevention and treatment of vitamin deficiencies, asthenia, aid resistance.
DOSAGE: I.M or S.C administration
- Bovine: 10 - 20 ml/ head.
- Calves, horse: 5 - 10 ml/ head.
- Sheep: 2 - 3 ml/ head.
- Pigs: 10 ml/ head.
- Piglets (10 kgs): 1 ml/ head.
- Weaners (10 - 30 kgs): 1 - 3 ml/ head.

COMPOSITION: 1 liter contains
Calcium gluconate.....100,000 mg
Vitamin D₃.....1,000,000 IU
Magnesium (chloride).....5,000 mg
INDICATIONS:
Supplement Calcium and essential substances for bone. Treatment of paralysis. Improvement of egg quality.

DOSAGE:
Poultry:
- Layer: 1 liter/ 1,000 liters of water, once per week.
- Broiler: 1 - 2 liter/ 1,000 liters of water, once per week.
- Cattle: 1 liter/ 1,200 litera of water.

CALCI-PLUS

500 ml, 1 liter



CALYP AZ

500 ml, 1 liter



COMPOSITION: 1 liter contains
Menthol.....20,000 mcg
Eucalyptol oil.....1,000 mg
Biotine.....5,000 mcg
INDICATION:
Antibacterial effect in the respiratory tract, decongestant and stimulate the local immune response system.

DOSAGE:
P.O. administration.
- Poultry: 1 ml/ 5 liters of drinking water.
- Small livestock: 10 - 20 ml/ animal.
- Large livestock: 20 - 40 ml/ animal.

COMPOSITION: 100 ml contains
Vitamin E.....10,000 mg
Sodium selenite.....100 mg
INDICATION:
- Vitamin E deficiencies
- Prevention of iron - toxication after administration of iron.

DOSAGE:
- Calves, goat, sheep: 1 ml/ 40 kgs B.W.
- Cattle: 1 ml/ 80 kgs B.W.
- Poultry: 1 liter/ 4,000 liters of drinking water.
- Chicks (<21 days): 1 liter/ 1,000 - 1,500 liters of drinking water.

E-SEL LIQUID

500 ml, 1 liter



▶ Oral solution

GALYVET

500 ml, 1 liter



COMPOSITION: 100 ml contains
 Alicin.....20%
DOSAGE:
 1 ml/ 2 liters of drinking water or 1 ml/ 10 - 15
 kgs B.W.

INDICATIONS:
 Boosting of intestine motility and enzymes
 secretion, stimulating appetite, increasing
 productivity. Detoxicate and keep healthy.

COMPOSITION: 1 liter contains
 Vitamin A.....500,000 IU
 Vitamin E.....5 g
 Selenium yeast.....800 mg
 Zinc sulphate.....10 g
 Beta glucan (yeast).....50 g
 Mannan oligosaccharide.....50 g
 Lectin.....50 g
 Thyme oil.....50 g

INDICATION
 Enhance growth rate, reduce the impact of
 stress in animals, increase the immune
 system and improve weight gain.
DOSAGE:
 - Poultry: 1 ml/ 4 liters of drinking water.
 - Cow & cattle: 1 ml/ 80 kgs B.W.
 - Sheep & goats: 1 ml/ 40 kgs B.W.

HEALTHVET

500 ml, 1 liter



HERBAVET

500 ml, 1 liter



COMPOSITION:
 Vitamins: Vitamin A 2,500,000 IU, Vitamin D₃ 500,000 IU,
 Vitamin E 3.75 g, Vitamin K₃ 250 mg, Pyridoxine 2 g,
 Thiamine 3.5 g, Riboflavine 4 g, Calcium Pantothenate 5 g,
 Cyanocobalamin 10 mg, Inositol 3 g.
 Amino acids: Methionine 16 g, Lysine 8 g.
 Inactive ingredient: Choline chlorine 40 mg, Proline 510 mg,
 Glycine 575 mg, Alanine 975 mg, Cystine 150 mg, Valine
 1.1 g, Leucine 1.5 g, Threonine 0.5 g, Tryptophane 75 mg,
 Histidine 0.9g, Arginine 490 mg, Aspartic acid 1.54 g, Serine
 680 mg, Glutamic acid 1.16 g, Isoleucine 1.25 g, Tyrosine
 340 mg, Phenylalanine 810 mg, Biotin 2 mg.

INDICATIONS: Treatment and prevention of vitamins
 and amino acids deficiency. Stimulate immune
 system, anti-stress on livestock and poultry.
DOSAGE:
 - General: 1 ml/ 4 - 5 liters per drinking water
 - Young poultry at brooding period: 1 ml/ 1.5 - 2 liters of
 drinking water.
 - Layer and broiler: 1 ml/ 2 - 3 liters of drinking water or
 1 ml/ 10 - 15 kgs B.W.
 - Finishing pig: 1 ml/ 2 - 3 liters of drinking water or 1 ml/
 20 - 30 kgs B.W.
 - Buffalos, cattle, goats, sheep: 1 ml/ 30 - 50 kgs B.W.

COMPOSITION: 1,000 ml contains
 Bromhexin HCl.....10,000 mg
DOSAGE:
 Mix with water: 1 - 2 ml/ 3 liters of drinking
 water.

INDICATIONS:
 Treatment of respiratory disorder such as:
 cough, pneumonia, chronic respiratory
 infection, bronchitis, pharyngitis, sinusitis.
 Expectorant.

HEXIN

500 ml, 1 liter



MEPHO

500 ml, 1 liter



COMPOSITIONS: 1 ml contains
 Menthol.....16 mg
 Camphor.....33 mg
DOSAGE:
 Clean and foment the affected area with hot
 water, then rub well with the MEPHO and
 cover (in necessary case).

INDICATIONS:
 - Treatment of myositis, bursitis, sprain
 tendons, tendovaginitis, tendinitis,
 lameness.
 - Relief of muscular soreness, aches and
 stiffness; and arthritic pains due to over-
 exertion and competitive activity.

COMPOSITION: 1 liter contains
 Fermented fungi essence (Saccharomyces).....35%
 Citric acid (min).....7%
 Lactic acid (min).....3%
 Formic acid (min).....2%
 Methionine (min).....40 g
 Lysine (min).....40 g
 Vitamin B₁ (min).....1,000 mg
 Vitamin B₂ (min).....1,000 mg
 Vitamin B₅ (min).....500 mg
 Vitamin B₆ (min).....500 mg

Vitamin PP (min).....1,000 mg
 Vitamin C (min).....2,000 mg
 Inositol (min).....2,000 mg
 ATP-Adenosine triphosphate (min).....2,000 mg
INDICATIONS:
 - Detoxication toxic out of the body.
 - Support energy resources to the body
DOSAGE:
 - Poultry: 1 ml/ 2 - 3 liters of drinking water or 100 ml/
 800 - 1,000 kgs B.W.
 - Cattle, horse: 1 ml/ 3 - 4 liters of drinking water or
 100 ml/ 1,000 - 1,200 kgs B.W.

MINOCID

500 ml, 1 liter





PHOCIUM

500 ml, 1 liter



COMPOSITION: 1 liter contains

Phosphorus.....	71 g
Calcium.....	2.9 g
Magnesium.....	0.8 g
Sodium.....	0.8 g
Manganese.....	1.6 g
Zinc.....	3.6 g
Copper.....	2.8 g
Cobalt.....	0.75 g

INDICATION:

Prevention of bone problem, improvement of egg shell.

DOSAGE:

1 ml/ 1 - 2 liters of drinking water.

COMPOSITION: 1 liter contains

Sorbitol.....	530 g
Methionine.....	20 g
Lysine.....	20 g
Betaine.....	50 g

DOSAGE:

- Piglets & dogs: 2 - 3 ml/ head.
- Swine: 5 ml/ head.
- Cattle, sheep, goats: 5 - 10 ml/ head.
- Poultry: 1 - 2 ml/ 1 liter of water.

INDICATION:

Enhance liver detoxification, prevent indigestion, feed poisoning and constipation. Stimulate appetite and weight gain.

PROLIVER

500 ml, 1 liter



SOBICAR

500 ml, 1 liter



COMPOSITION: Each ml contains

Sodium Bicarbonate.....	100 mg
-------------------------	--------

INDICATION:

Treatment of digestive disorders, indigestion, acidosis and increases the level of fat in milk production.

DOSAGE:

- Poultry: 1 ml/ animal/ day.
- Cows: 40 ml/ animal/ day.
- Horse: 25 ml/ animal/ day.
- Goats: 15 ml/ animal/ day.
- Sheep, camels & calves: 20 ml/ animal/ day.

COMPOSITIONS: 1 liter contains

Propionic acid.....	60 ml
Citric acid.....	10 g
Choline chloride.....	500 mg
Ferric ammonium citrate.....	200 mg
Zinc chloride.....	500 mg
Protein hydrolysate.....	100 mg
Copper sulphate.....	300 mg
Vitamin B ₂	75 mg
Vitamin B ₆	200 mg

Vitamin B ₁₂	2 mg
Magnesium chloride.....	500 mg
Yeast extract.....	2%

INDICATIONS:

Acts well on toxin and moulds. Increase the quality of water by reducing pH.

DOSAGE:

1 ml/ 1 liter of water 5 - 7 days every month in water 250 - 500 ml/ metric ton of feed continuously

TOXYLIQ

500 ml, 1 liter



VITAMI COMPLEX

50 ml, 100 ml



COMPOSITIONS: 100 ml contains

Vitamin A.....	50,000 IU
Vitamin D ₃	10,000 IU
Vitamin E.....	10 mg
Vitamin B ₁	2.5 mg
Vitamin B ₆	1.5 mg
Nicotinamide.....	17.5 mg
Vitamin C.....	25 mg
Vitamin K ₃	1.5 mg

INDICATIONS:

Stimulate appetite on poultry, fast growth.

DOSAGE:

Pour directly or mix with water
Poultry: 2 - 5ml/ 1 liter of water

COMPOSITION: 1 kg contains

Humidity.....	12%
Biotin.....	6,000 mcg
ZnSO ₄	4,500 - 5,500 mg
Calci Pantothenate.....	2,500 mg
Methionine.....	60,000 mg
Folic acid.....	300 mg
Vitamin A.....	2,000,000 IU
Vitamin D ₃	1,000,000 IU
Vitamin E.....	1,500 mg

Vitamin B ₁	3,000 mg
Vitamin B ₂	2,000 mg
Vitamin B ₆	3,000 mg
Niacin.....	3,000 mg

INDICATIONS:

Supply acid amines, vitamins, minerals to prevent cannibalism, hair loss

DOSAGE: For oral administration

1 - 2 g/ 1 liter of drinking water or 10 g/ 10 kgs B.W, daily using 100 g/ 30 kgs feed



ACIVITA

500 g, 1 kg



Powder

ADE THIAM

500 g, 1 kg



COMPOSITION: 1 kg contains
 Vitamin A (min).....1,000,000 IU
 Vitamin D₃ (min).....200,000 IU
 Vitamin E (min).....1,000 mg
 Methionine (min).....1,500 mg
 Vitamin B₁ (min).....200 mg
 Vitamin B₆ (min).....100 mg
 Vitamin PP (min).....200 mg
Bacillus subtilis (min).....10⁸ CFU
 Zinc sulfate (min - max).....720 - 880 mg

Iron sulfate (min - max).....360 - 440 mg
 Copper sulfate (min - max).....90 - 110 mg
 Milky powder q.s.....1 kg

INDICATIONS:

- Supply of essential microorganism for digestive tract, prevention of digestive disorder.
 - Provide necessary nutrition for stunted swine, grow fast, prevent diarrhea.

DOSAGE: For oral administration

Mix with feed: 1 kg/100 kgs of feed or 2 g/ 1 liters of water.

COMPOSITION: 1 kg contains

Vitamin C.....20%

INDICATIONS:

Increase resistance, anti-stress caused by transportation or climate change, anti - depression.

DOSAGE:

- Livestock: 100 g/ 120 kgs of feed or 100 g/ 280 liters of water.

- Poultry: 100 g/ 150 - 200 liters of water.

ASCOBIC-WS

500 g, 1 kg



BLOOVET

500 g, 1 kg



COMPOSITION: 50 g contains

Iron.....600 mg
 Zinc.....24 mg
 Manganese.....9.6 mg
 Copper.....24 mg
 Vitamin B₁₂.....1,200 mcg
 Vitamin B₆.....9.6 mg
 Vitamin B₁.....25 mg
 Folic acid.....30 mg
 Methionine.....1,500 mg

Vitamin C.....600 mg
 Vitamin K₃.....60 mg
 Vitamin E.....577 mg

INDICATIONS:

For good blood maintenance a boost before and after racing.

DOSAGE:

Mix in feed or drench
 - Young camel: 25 g/ head.
 - Adult animal: 50 g/ head.

COMPOSITION: 1 kg contains

Vitamin A (min).....3,500,000 IU

Biotin (min).....4,000 mcg

Zinc sulfate (min-max).....90,000 - 110,000 mg

INDICATIONS:

Improve skin, hair, hoof condition in cattle, goats, sheep, horse

DOSAGE:

Cattle, goats, sheep: 2.5 - 3 kg/ ton of mixed feed or 1 - 2 g/ 50 kgs B.W, for 3 - 5 days.

DETOXVET

500 g, 1 kg



DIGEST-TONIC

500 g, 1 kg



COMPOSITION: 101 g contains

Magnesium carbonate.....90 g

Sodium carbonate.....10 g

Nux vomica.....1 g

INDICATION:

Treatment of cases of indigestion in animals. Stimulate appetite, ruminal tonic and detoxicant.

DOSAGE:

Large animals: 100 g/ liter of drinking water.

Small animals: 40 g/ ½ liter of drinking water.

COMPOSITION: 1 kg contains

Vitamin A (min).....10,000,000 IU

Vitamin D₃ (min).....500,000 IU

Vitamin E (min).....2,000 IU

Vitamin B₁ (min).....1,000 mg

Vitamin B₂ (min).....1,500 mg

Vitamin B₃ (min).....3,000 mg

Vitamin B₆ (min).....1,000 mg

Vitamin B₁₂ (min).....10,000 mcg

Vitamin K₃ (min).....500 mg

Niacinamide (min).....1,000 mg

Biotin (min).....100 mcg

Folic acid (min).....100 mg

Sodium chloride (min - max).....9,000 - 11,000 mg

Potassium chloride (min - max).....4,500 - 5,500 mg

Excipient (ginseng) q.s.....1 kg

INDICATIONS:

- Supporting for immunology system, prevention of stuntedness, increasing growth rate.
 - High health recovery after treatment.

DOSAGE: For oral administration

Mix with feed or drinking water
 - Prevention of stuntedness, poor growth: 20 g/ 100 animals/ day or 20 g/ 20 liters of drinking water.

- Stimulation of growth rate, health recovery: 10 g/ 100 animals/day or 10 g/ 20 liters of drinking water.

- Prevention and treatment of Gumboro: 20 g/ 5 liters of drinking water.

GINSENG

(Granular nutrient)

500 g, 1 kg





HAEZET

500 g, 1 kg



COMPOSITION: 1 kg contains
 Vitamin K₃.....10 g
INDICATIONS:
 Treatment of hemorrhage due to vitamin K deficiency. Bleeding after giving birth.

DOSAGE:
 For oral administration
 - Cattle: 1 g/ 2.5 liters of water.
 - Poultry: 1 g/ 2 liters of water.

COMPOSITION: 200 g contains
 Humidity (max).....13%
 Glucose (min).....66 g
 Lysine (min).....10 g
 Vitamin C (min).....20 g
 Vitamin K₃ (min).....9 g
 Na+ (min - max).....3.6 - 4.4 g
 K+ (min - max).....2.4 - 2.8 g
 Calcium gluconate (min - max).....18 - 22 g
 Grain of sand (max).....0.5%

INDICATIONS:
 - Supplement vitamins, amino acids and electrolytes for poultry, cattle detoxification, disease resistance
 - Increase absorption and digestion, growth and metabolism, increase the survival rate
DOSAGE:
 Mix with feed or water
 1 g/ 2 - 3 liters of drinking water

HEALBALAN

500 g, 1 kg



HEASUPPORT

500 g, 1 kg



COMPOSITION: 1 kg contains
 Vitamin A.....12,000,000 IU
 Vitamin D₃.....4,000,000 IU
 Vitamin E.....25,000 mg
 Vitamin B₂.....7,000 mg
 Vitamin PP.....50,000 mg
 Vitamin C.....50,000 mg
 Methionine.....2,300 mg
Bacillus subtilis.....2x 10⁹ CFU
 Calcium.....10,000 mg

INDICATIONS:
 - Vitamin and amino acid supplement. Prevention of disease caused by vitamin deficiency.
 - Prevention of stress caused by heating. Stimulation growth rate.
DOSAGE:
 - Poultry: 250 - 500 g/ ton of feed or 500 g/ 1,000 liters of drinking water.
 - Cattle, buffalos, horse: 500 g/ 1.5 - 2 tons of feed or 500 g/1,000 liters of drinking water.

COMPOSITION: 1kg contains
 Methionine (min).....30,000 mg
 Lyzine (min).....5,000 mg
 ZnSO₄ (min - max).....1,800 - 2,200 mg
 Vitamin B₁ (min).....500 mg
 CaCO₃ (min - max).....787 - 962 g

INDICATIONS:
 Increasing milk secretion in sows and dairy cows.
DOSAGE:
 For oral administration
 The product for sows, milk cows: 10 g/ time, twice per day.

MILK ENHAN

500 g, 1 kg



PHEROL WS

500 g, 1 kg



COMPOSITION: 1 kg contains
 Vitamin E.....1,200 IU
INDICATIONS:
 Treatment and prevention Vitamin E deficiency in pigs, poultry, cattle, goats.

DOSAGE:
 Poultry: 5 g/ 3 - 5 liters of drinking water
 Pigs, cattle, buffalos: 5 g/ 50 kgs B.W.
 Or mix 100 g/ 50 - 100 kgs feed

COMPOSITION:
 Consist of *Bacillus subtilis* spp., *Lactobacillus* spp., *Saccharomyces* spp., digestive enzyme α- Amylase, Protease.
 1 kg contains:
Bacillus subtilis.....1 x 10⁹ CFU
Lactobacillus acidophilus.....1 x 10⁷ CFU
Saccharomyces cerevisiae.....1 x 10⁷ CFU
 Vitamin B₁.....1,000 mg
 Excipient (Glucose, Garlic,...) q.s.....1 kg

INDICATIONS: The product supplies alive profitable bacterium for digestive system on livestock and poultry. Prevention of digestive disorders. Stimulation of digestive activity, increasing growth rate. Prevention of diarrhea caused by abnormal food and water. Should be given after antibiotics administration to recover digestive function.
DOSAGE: Dilute into water or mix with feed
 - Poultry: + Against: 1 g/ 1 litter of drinking water or 1 g/ 10 kgs B.W or 1 kg/ 1 ton of feed.
 + Prevention: ½ of treatment dose.
 - Cattle:
 + Against: 1 g/ 1 liter of drinking water or 2 - 3 g/ cattle/ twice per day.
 + Prevention: ½ treatment dose

PROBI-ALLY

(Granular nutrient)

500 g, 1 kg



Powder

PROBIZ

500 g, 1 kg



COMPOSITON: 1 kg contains
Bacillus subtilis (min).....10⁸ CFU
 Methionine (min).....16,000 mg
 Thiamin nitrate (min).....15,000 mg
 Sodium chloride (min - max).....1.98 - 2.42%
 Potassium chloride (min - max).....1.17 - 1.43%
 Butaphosphan (min).....20,000 mg
 Sorbic acid (min).....100 mg
 Citric acid (min).....150 mg
 Zinc sulphate, ferrous sulphate, magnesium sulphate (min - max).....4.770 - 5.830 mg
 Natural and artificial flavours (min).....3,800 mg

INDICATION:
 Anti - stress caused by vaccination, transportation, temperature.
DOSAGE
 1 g/ 1 - 2 liter of water or 1 - 2 kg/ ton of feed.

COMPOSITION: 1 kg contains
Bacillus subtilis.....10⁹ CFU
Saccharomyces cerevisiae.....10⁶ CFU
Lactobacillus acidophilus.....10⁶ CFU
 Sodium chloride......3%
 Potassium chloride.....2.5%
DOSAGE:
 - Cattle: 250 - 500 g/ 1 ton of feed.
 - Poultry: 100 g/ 100 litres of water.

INDICATIONS:
 - Stimulate digestion, appetite and food intake and keep healthy.
 - Balance intestinal bacteria, prevent intestinal infection like diarrhea caused by *E.coli*, *Salmonellosis*, necrotizing enterocolitis in poultry.
 - Prevention and treatment of diarrhea in piglets, indigestion, digestive disorder caused by antibiotic using.

PROTIC

500 g, 1 kg



RACE COMPE

500 g, 1 kg



COMPOSITION: 25 g contains
 DI-lipoic acid.....1,200 mg
 Vitamin B₁.....500 mg
 Pantothenic acid.....450 mg
 Riboflavin.....35 mg
 Niacin.....35 mg
INDICATION: Is a pre-performance supplement for horse during intense activity when need for energy is greatest.

DOSAGE:
 For support under stressful condition, feed 25 g daily for 2 days. Thereafter feed 12 - 25 g as needed. For stress or performance, give 25 g the evening before an event and 25 g two to three hours prior to the event

COMPOSITION: 100 g contains
 Sodium chloride.....6.2 g
 Potassium chloride.....4 g
 Sodium bicarbonate.....3 g
 Sodium citrate.....6.5 g
 Calcium lactate.....1.7 g
 Magnesium sulphate.....1 g
 Vitamin C.....600 mg
 Dextrose.....77 g

INDICATION:
 - In summer to avoid dehydration and heat stroke.
 - As an aid in diarrhea and dysentery.
 - To maintain correct osmotic pressure and precise acid base equilibrium.
DOSAGE: Administration through drinking water.
 - Cows, buffaloes: 60 g daily.
 - Calves, sheep and goat: 30 g daily.
 - Dogs: 5 - 10 g per day.
 - Horse and camels: + Light work: 30 g per day.
 + Moderate work 60 g per day.
 + Intense work 90 g per day.

REHYDRATE

500 g, 1 kg



RINEVET

500 g, 1 kg



COMPOSITION: 1 g contains
 Ammonium chloride.....650 mg
 DL - Methionine.....100 mg
 Vitamin A.....250 IU
 Vitamin C.....100 mg
DOSAGE:
 Poultry: 1 g/ liter of drinking water or 2 kg/ ton of feed

INDICATION:
 RINEVET is a diuretic, supportive treatment of urinary tract infections. Enhance the hepatic function and regulation of cholesterol.

COMPOSITION: 100g contains
 Butafosfan.....10,000 mg
 Cyanocobalamin.....3,000 mcg
 Biotin.....12,000 mcg
 ZnSO₄.....2,000 mg
 Vitamin A.....1,000,000 IU
 Vitamin D₃.....400,000 IU
 Vitamin E.....2,000 IU
INDICATIONS: Aid absorption and transformation of calcium, phosphor, prevent stunt. Stimulate hair growth. Prevent cannibalism, jaundice.

DOSAGE:
 For oral administration
 - Poultry:
 + 1 - 40 days of age: 100 g/ 400 liters of water
 + 40 - 90 days of age: 100 g/ 300 liters of water or 100 g/ 80 kg of feed.
 + 90 days of age onward: 100 g/ 400 liters of water or 100 g/ 150 kg of feed.
 - Livestock: 100 g/ 600 liters of water or 100 g/ 300 kgs of feed.

SORBIVET

500 g, 1 kg



Powder



SUPPLEVET

500 g, 1 kg



COMPOSITION: 1 kg contains
 Vitamin A.....8,000,000 IU
 Vitamin D₃.....1,500,000 IU
 Vitamin E.....1,000 mg
 Vitamin B₁.....500 mg
 Vitamin B₂.....500 mg
 Vitamin B₆.....200 mg
 Vitamin B₁₂.....8,000 mcg
 Folic acid.....50 mg
 Ca-D pantothenate.....4,000 mg
 Vitamin K₃.....2,000 mg
 Nicotinamide.....6,000 mg
 Manganese.....400 mg
 Zinc.....150 mg
 Iron.....500 mg
 Copper.....40 mg
 Cobalt.....10 mg
 L-Lysine.....61 mg
 DL- Methionine.....13 mg
 Glycine.....382 mg
 Alanine.....146 mg
 Arginine.....177 mg

Cysteine.....10 mg
 Leucine.....48 mg
 Phenylalanine.....40 mg
 Valine.....40 mg
 Glutamic acid.....166 mg
INDICATIONS:
 Supplementation with essential micro nutrients.
DOSAGE:
 1 kg/ 2,000 liters of drinking water.

COMPOSITION: 1 kg contains

Humidity.....12%
 Vitamin A.....1,000,000 IU
 Vitamin D₃.....100,000 IU
 Vitamin E.....4,000 IU
 KCl.....42,500 mg
 NaCl.....17,000 mg
 Vitamin PP.....5,000 mg
 Vitamin B₆.....1,200 mg
 Biotin.....500 mcg
 Vitamin B₁₂.....6,000 mcg

INDICATIONS:

Enhanced growth and resistance, increase the survival rate, egg production, preventing stress.
DOSAGE:
 Mixed in food or water
 - Calves, goats, sheep: 1 g/ 4 kgs of feed or 1 g/ 4 liters of water.
 - Poultry: 1 g/ 4 kgs of feed or 1 g/ 4 liters of water.

TAVILEX

500 g, 1 kg



TROLYT AIZ

500 g, 1 kg



COMPOSITION: 100 g contains
 Sodium bicarbonate (min - max)....36,000 - 44,000 mg
 Sodium chloride (min - max).....23,400 - 28,600 mg
 Potassium chloride (min - max).....4,500 - 5,500 mg
 Dextrose monohydrate.....29,000 mg

INDICATION:

Treatment of dehydration, disorder of electrolytes caused by fever or diarrhea and anti stress.
DOSAGE: P.O. administration
 1 - 2 g/ liter of drinking water or 4 - 5 g/ 1 kg of feed.

COMPOSITIONS: Consists of vitamins, acid amines, mineral, profitable microorganism as: *Bacillus subtilis*, *Lactobacillus acidophilus*, *Saccharomyces cerevisiae*, *Aspergillus oryzae*...

1 kg contains
 Phosphorus (min - max).....42.5 - 57.5 g
 Calcium (min - max).....229.5 - 310.5 g
 Vitamin A (min).....500,000 IU
 Vitamin D₃ (min).....60,000 IU
 Vitamin E (min).....500 IU
 Biotin (min).....4 mcg
 Cobalt sulfate (min - max).....18 - 22 mg

Iodine (min).....20 mg
 FeSO₄ (min - max).....9,000 - 11,000 mg
 CuSO₄ (min - max).....4,500 - 5,500 mg
 ZnSO₄ (min - max).....18,000 - 22,000 mg
 MnSO₄ (min - max).....18,000 - 22,000 mg
INDICATIONS: Supplement of vitamins, mineral, digestive enzymes for bone formation, prevention of malnutrition, paralysis after giving birth on dairy cow. Stimulation of growth, increasing productivity and milk quality.

DOSAGE:
 - Dairy cow: 20 g/ cow or 1 kg/ 150 kgs feed.
 - Beef: 1 kg/ 400 kgs feed.
 - Calves, goat, sheep: 1 kg/ 300 kgs feed.

VIRALMIN

500 g, 1 kg



VITA-CA

500 g, 1 kg



COMPOSITION: Each 1 g contains
 Calcium.....42 mg
 Phosphorus.....15 mg
 Vitamin B₁₂.....10 mcg
 Vitamin C.....6 mg
 Vitamin D₃.....400 IU
 Citric Acid.....27 mg

INDICATION:

Prevention & treatment of rickets, osteoporosis & anemia. Formation of skeletal structure & egg shell.
DOSAGE:
 - Chicks: 1 g/ 1 liter of water.
 - Broiler: 1 g - 2 g/ 1 liter of water.
 - Layer: 2 - 4 g/ 1 liter of water.

COMPOSITION: 100 g contains

Vitamin A.....2,500,000 IU
 Vitamin D₃.....200,000 IU
 Vitamin E.....1,200 mg

INDICATIONS:

- Prevent and treat deficiency of Vitamin A, D, E.
 - Prevent abortion due to Vitamin deficiencies.
 - Supplement vitamins for poultry, cattle.

DOSAGE:

- Poultry: 500 g/ 1,200 liters of water.
 - Swine, goats & cattle: 500 g/ 2,400 liters of water or 500 g/ 1,000 kgs of feed.

VITASUPLY

500 g, 1 kg



ETD

50 ml, 100 ml



COMPOSITION: 1 ml contains
Estradiol benzoate.....2 mg

INDICATION:
Supply estrogen for female livestock, stimulate oestrus and ovulation, increase efficiency of oosperm, and the number of embryo.

DOSAGE:
For I.M administration
- Horse: 5 - 10 ml.
- Sheep: 0.6 - 1 ml.
- Cattle: 2.5 - 10 ml.
- Dog: 0.1 - 0.2 ml.

COMPOSITION: 10 ml contains
Estradiol benzoate.....50 mg
Vitamin A.....125,000 IU
Vitamin E.....50 mg

INDICATION
Stimulate estrus and ovulation, increase efficiency of oosperm and the number of embryo. Treatment of hysteritis, uterus pyogenesis and prevent deficiency of vitamin A, E.

DOSAGE:
S.C administration
- Horses: 2 - 4 ml.
- Sheep, goats: 0.2 - 0.4 ml.
- Cattle: 1 - 4 ml.
- Dogs: 0.04 - 0.1 ml.

ETD VIT

50 ml, 100 ml



TOXYL

50 ml, 100 ml



COMPOSITION: Each 100 ml contains
Oxytocin.....1,000 IU

INDICATION:
For treatment of weakness, pain in the course of delivery, lactation failure in sows, uterus atonia, retention of the placenta, postpartum hemorrhage, prolapse of the uterus, atonia of the intestine and the urinary bladder, postpartum agalactia and torsion of gravid uterus.

DOSAGE:
I.M, S.C administration
- Mares, cows: 4 - 5 ml per animal.
- Ewes, goats: 1 - 3 ml per animal.
- Sows: 2 - 4 ml per animal.
- Bitch: 0.5 - 1 ml per animal.
- Cats: 0.3 - 0.5 ml per animal.





ANAZIN.3000

50 ml, 100 ml



COMPOSITION: 100 ml contains
 Analgin.....30 g
INDICATIONS:
 - Pain relief, rheumatism prevention, antipyretic.
 - Treatments of rheumatism, intestinal spasm, intestinal peristalsis.

DOSAGE:
 I.V, I.M administration
 - Bovine, horses: 1 ml/ 15 - 20 kgs B.W.
 - Goats, sheep, swine: 1 ml/ 10 - 15 kgs B.W.

COMPOSITION: 1 ml contains
 Phenylbutazone..... 200 mg
DOSAGE: I.V. administration
 Horses: 1 ml/ 45 kgs, with a maximum of 10 ml.

INDICATIONS:
 Analgesic, reduce fever, anti-inflammatory and treatment of musculo-skeletal disorders (e.g. osteoarthritis, bursitis, lameness,...).

BUTAZONE-Z

50 ml, 100 ml



CORHENYL

50 ml, 100 ml



COMPOSITION: 100 ml contains
 Phenylbutazone.....18 g
 Dexamethasone sodium phosphate....35 mg
 Benzyl alcohol.....1 ml
 Disodium edetate.....4 mg
INDICATION:
 Anti-inflammation, analgesic, antipyretic.

DOSAGE:
 I.V. and I.M. administration
 - Horse and camel: 4 ml/ 100 kgs B.W.
 - Dogs: 0.5 ml/ 10 kgs B.W.

COMPOSITION: 100 ml contains
 Dexamethasone sodium phosphate.....5 mg
 Ammonium Chloride.....150 mg
INDICATION:
 High effects on anti-inflammation and anti - allergic, reduce swelling, anti - congestion and enhancing resistance during exercise.

DOSAGE:
 Slow I.V or I.M administration
 - Prevention: 10 ml/ animal.
 - Treatment: 20 ml/ animal.

CORNIUM

50 ml, 100 ml



CORTYL VET

50 ml, 100 ml



COMPOSITIONS: 100 ml contains
 Dexamethasone sodium phosphate...100 mg
INDICATIONS:
 Anti - inflammation and anti - allergic agent in livestock, poultry, dogs and cats.

DOSAGE:
 I.M administration
 - Big animals: 1 ml/ 20 kgs B.W .
 - Young animal: 1 ml/ 10 kgs B.W .
 - Poultry: 1 ml/ 5 kgs B.W.

FORMULATION: 1 ml contains
 Triamcinolone acetone.....2 mg
INDICATIONS:
 Treatment of inflammations caused by infection, allergies, traumatism.

DOSAGES: Intramuscular or subcutaneous administration
 - Equine: 6 - 10 ml/ head.
 - Dairy cows: 10 ml/ head.
 - Dogs and cats: 0.05 - 0.1 ml/ kg B.W.

CYNOLON

50 ml, 100 ml



▶ **Injectable solution**



COMPOSITION: 1 ml contains
 Prednisolone.....7.5 mg
 Dexamethasone sodium phosphate...2.5 mg
INDICATIONS:
 Arthritis, synovitis, inflammation of the tendon sheaths, strains, itching, eczema, dermatitis.

DOSAGE:
 I.V, I.M and S.C administration
 - Horses & camels: 3 ml/ 100 kgs B.W.
 - Poultry: 3 ml/ 25 kgs B.W.

COMPOSITION: 1 ml contains
 Diclofenac sodium.....100 mg
 Solvent q.s.....1 ml
INDICATIONS:
 Anti-inflammatory, analgesic and anti-pyretic.

DOSAGE:
 - Intramuscular injection for 1 - 3 consecutive days.
 - Pigs, cattle, camels, horse & goats: 1 ml/ 40 kgs B.W/ day.
 - Dogs & cats: 0.25 ml/ 15 kgs B.W/ day.



COMPOSITIONS: 1ml contains
 Hydrochlorothiazide.....50 mg
 Dexamethasone sodium phosphate..0.5 mg
INDICATIONS:
 For anti-inflammation in livestock.

DOSAGE: I.V, I.M, S.C
 Cattle and adult horses.
 - Preventive: 10 ml daily.
 - Treatment:
 + Congestion & mild edema: 10 ml.
 + Congestion & severe edema: 20 ml.
 Foals and calves: 2 ml/ 40 - 50 kgs of B.W.

COMPOSITIONS: 100 ml contains
 Bromhexin HCl.....300 mg
 Dipyrone.....20,000 mg
 Dexamethasone sodium phosphate...50 mg

INDICATIONS:
 Anti - inflammation, pain relief, antipyretic, expectorant. Treatment of cough, dyspnea
DOSAGE: I.M administration
 1 ml/ 10 kgs B.W.



COMPOSITION: 100 ml contains
 Analgin.....25 g
 Vitamin C.....10 g
INDICATIONS:
 Antipyretic, pain and strengthened resistance.

DOSAGE:
 - Intramuscular administration
 - Swine, goats, sheep, calves: 1 ml/ 10 kgs BW.
 - Bovine, sows, boars: 1 ml/ 15 - 20 kgs B.W.

COMPOSITION: 1 ml contains
 Flumethasone.....0.25 mg
 Dexamethasone.....0.25 mg
 Vitamin C.....9 mg
INDICATION:
 - Special formula for high performance competition in racing animal.
 - High anti-inflammatory effect, anti-allergic, anti stress and enhance resistance.

DOSAGE:
 I.M. administration
 - Large livestock: 4 - 8 ml.
 - Dogs:
 + Small and medium: 0.25 - 0.5 ml.
 + Large: 0.5 - 1 ml.
 - Cats: 0.1 - 0.3 ml.
 - Large livestock: 4 - 8 ml.



FLUMEA

50 ml, 100 ml



COMPOSITION: 1 ml contains
 Flumethasone.....0.25 mg
 Vitamin C..... 9 mg
INDICATION:
 Supportive therapy in milk fever. Treatment of various rheumatic, musculoskeletal inflammation, mastitis, metritis, etc., allergic. Enhance resistance.

DOSAGE:
 I.M, I.V, S.C administration
 - Bovine and equine: 5 - 10 ml.
 - Canine:
 + Small and medium: 0.4 - 1 ml.
 + Large: 1 - 2 ml.
 - Feline: 0.2 - 0.5 ml.,

COMPOSITION: 100 ml contains
 Ketoprofen.....10,000 mg
INDICATIONS:
 Inflammatory and painful of the bones, joints and muscle-skeletal systems in livestock

DOSAGE:
 - Pigs, bovine, goats, sheep: intramuscular or intravenous injection 1 ml/ 33 kgs B.W.
 - Horses: intravenous injection 2,2 ml/ 100 kgs B.W.
 - Dogs, cats: intravenous injection 0,1 ml/ 10 kgs B.W.

KEFEN

50 ml, 100 ml



NIXIN-F

50 ml, 100 ml



COMPOSITION: 100ml contains
 Flunixin meglumine.....6 g
INDICATION:
 Livestock: treatments of inflammatory, respiratory diseases, pneumonia, mastitis, metritis, nephritis, fever.

DOSAGE:
 - Swine, bovines, goats, sheep:
 + I.M injection: 2 ml/ 45 kgs B.W.
 + Musculoskeletal inflammation and disorder: 1 ml/ 45 kgs B.W for I.V injection
 - Horses: 1 ml/ 45 kgs B.W for I.V injection.

COMPOSITION: 1 ml contains
 Prednisolone sodium phosphate.....25 mg
DOSAGE:
 I.M, I.V admistration
 General dose: 5 ml/ head.

INDICATIONS:
 - Relief of inflammation and clinical parameters associated with recurrent pathway obstruction.
 - Respiratory in horses, in combination with environmental control.

SOLINE-Z

50 ml, 100 ml



TRYFASIN

50 ml, 100 ml



COMPOSITIONS: 100 ml contains
 α - Chymotrypsin.....400 mg
 Trypsin.....100 mg
INDICATION:
 Anti - inflammation, reduce edema, pain and swelling after surgery. Supportive treatment of mastitis, arthritis, asthma, bronchitis, pneumonia, sinusitis.

DOSAGE:
 Intramuscular administration
 - Poultry: 1 ml/ 10 - 15 kgs B.W.
 - Young livestock: 1 ml/ 20 - 30 kgs B.W.
 - Mature livestock: 3 - 6 ml/ 100 kgs B.W.

COMPOSITIONS: 100 ml contains
 α - Chymotrypsin.....400 mg
 Trypsin.....100 mg
INDICATION:
 Anti - inflammation, reduce edema, pain and swelling after surgery. Supportive treatment of mastitis, arthritis, asthma, bronchitis, pneumonia, sinusitis.

DOSAGE: For oral administration
 - Poultry: 1 ml/ 8 - 12 kgs B.W. or 1 ml/ 1 - 2 liters of drinking water.
 - Calves, lamb, goatling: 1 ml/ 20 - 30 kgs B.W.
 - Cattle, buffalos, sheep, goats: 3 - 6 ml/ 100 kgs B.W.
 Interval dose 12 - 24 hours if necessary.
 * Medicine should be used for 5 - 7 days since unboxing.

TRYPFA

500 ml, 1 liter



Powder

DEROZI-PR

500 g, 1 kg



COMPOSITION: 100 g contains
 Hydrochlorothiazide.....7.5 g
 Dexamethasone sodium phosphate...0.025 g
DOSAGE:
 40 g product on day 1.
 20 g product on day 2 and 3.

INDICATIONS:
 Livestock: Udder and perineal edema, lung edema, wound swelling after surgery and edema caused by allergy.

COMPOSITION: 100 g contains
 Analgin.....10,000 mg
 Vitamin C.....10,000 mg
INDICATIONS:
 Pain relief, antipyretic on acute infected cases.

DOSAGE:
 - Poultry: 100 g/ 70 - 100 liters of water or 100 g/ 70 - 80 kgs of feed.
 - Livestock: 100 g/ 100 - 130 liters of water or 100 g/ 200 kgs of feed.

METAMIC

500 g, 1 kg



PARA+C

500 g, 1 kg



COMPOSITION: 100 g contains
 Paracetamol.....10 g
 Vitamin C.....10 g
INDICATIONS:
 - Antipyretic, pain relief, influenza, resistance aid when climate changes.
 - Pain relief, antipyretic in acute infection.

DOSAGE:
 - Poultry: 100 g/ 80 - 100 liters of water or mix with 80 kgs of feed.
 - Cattle: 100 g/ 100 - 130 liters of water or mix with 100 kgs of feed.

COMPOSITION: 1 kg contains
 Aspirin (Acetylsalicylic acid).....500 g
INDICATIONS:
 - Anti-inflammation, pain relief, antipyretic.
 - Combination with antibiotic for treatment of infectious diseases, acute fever.
 - Safe for dogs, cats.

DOSAGE:
 Mix with feed for 3 - 5 days depend on cases
 - Large livestock: 0.3 - 0.5 g/ kg B.W/ day.
 - Small livestock, poultry: 0.2 - 0.25 g/ kg B.W/ day.

PYRIVET

500 g, 1 kg



SALIC

500 g, 1 kg



COMPOSITION: 100 g contains
 Acetyl salicylic acid.....10 g
 Ascorbic acid.....10 g
INDICATIONS:
 Analgesic and antipyretic cases of vaccination and transportation.

DOSAGE:
 Mix with drinking water
 1 g/ 2 liters drinking water
 Excessive heat: 1 g/ 1 liter drinking water





COLISTIN

50 ml, 100 ml



COMPOSITION: 100 ml contains
Colistin sulfate.....50,000,000 UI

INDICATIONS:
Treatment of respiratory and digestive disease, mastitis, metritis.

DOSAGE:
For I.M administration
- Swine, goats: 1 ml/ 7 - 10 kgs B.W.
- Bovine, horses: 1 ml/ 10 - 15 kgs B.W.
- Poultry: 2 ml/ 1 liter of water

COMPOSITION: 100 ml contains
Tylosin tartrate.....20 g
Colistin sulfate.....50,000,000 UI
Dexamethasone.....100 mg

INDICATIONS:
Treatment of acute pneumonia, pasteurellosis, enteritis, edema, diarrhea, arthritis, metritis, mastitis, etc.

DOSAGE:
I.M administration for 3 - 5 days
- Piglets, kids, lamb: 1 ml/ 10 kgs B.W.
- Swine, calves, goats, sheep: 1 ml/ 15 kgs B.W.
- Buffalos, cattle: 1 ml/ 20 - 30 kgs B.W.
- Poultry: 1 ml/ 5 - 7 kgs B.W.

COLYXA

50 ml, 100 ml



COSPIRA

50 ml, 100 ml



COMPOSITION: 100 ml contains
Spiramycin adipate.....32,000,000 IU
Colistin sulfate.....40,000,000 IU

INDICATION:
For treatment of respiratory and digestive disease in livestock and poultry.

DOSAGE:
I.M injection
- Livestock: 1 ml/ 10 - 12 kgs B.W per day.
- Poultry: 1 ml/ 6 - 8 kgs B.W per day.

COMPOSITION: 100 ml contains
Gentamycin sulfate.....6,000 mg
Colistin sulfate.....50,000,000 UI
Dexamethasone sodium phosphate.....200 mg

INDICATIONS:
Treat CRD, enteritis, *E.coli*, omphalitis, diarrhea, swollen head in ducks, pasteurellosis, salmonellosis and chronic sinusitis.

DOSAGE:
Intramuscular injection for 3 - 5 days.
- Poultry: 1 ml/ 5 - 10 kgs B.W.
- Cattle: 1 ml/ 20 kgs B.W.
- Swine, Goats & Sheep: 1 ml/ 10 - 15 kgs B.W.

DECOGEN

50 ml, 100 ml



DOF-SONE

50 ml, 100 ml



COMPOSITION: 100 ml contains
Doxycycline hyclate.....10 g
Florfenicol.....10 g

INDICATION:
Treatment of respiratory and digestive diseases. Against of mastitis - metritis and swollen head.

DOSAGE:
I.M administration
- Bovine, horses, swine: 1 ml/ 12 - 15 kgs B.W.
- Piglets, calves, goats, sheep: 1 ml/ 10 kgs B.W.

COMPOSITION: 100 ml contains
Spiramycin adipate.....20,000,000 IU

INDICATIONS:
Against of pneumonia, enteritis, salmonellosis, arthritis, mastitis, metritis, vaginitis, wound, genital tract infection.

DOSAGE:
I.M administration
- Cattle, horse: 1 ml/ 15 - 20 kgs B.W.
- Calves, goat, sheep: 1 ml/ 10 kgs B.W.

DYPATE

50 ml, 100 ml



▶ Injectable solution

FLOCOVET

50 ml, 100 ml



COMPOSITION: 1 ml contains
 Enrofloxacin base.....25 mg
 Colistin sulfate.....250,000 IU
 Bromhexine HCl.....0.5 mg
INDICATION:
 For treatment of intestinal infections,
 respiratory infection, urethritis, mastitis,
 metritis, arthritis, foot rot.

DOSAGE:
 For intramuscular administration for 3 - 5 days.
 - Cattle, sheep, goats: 1 ml/ 10 - 12 kgs B.W.
 - Poultry, dogs, cats: 1 ml/ 8 - 10 kgs B.W.

COMPOSITION: 100 ml contains
 Florphenicol.....40 g
INDICATIONS:
 Treatment of respiratory infection.

DOSAGE:
 I.M injection
 Livestock: 1 ml/ 20 - 30 kgs B.W.

FLORJECT

50 ml, 100 ml



FLOTYL

50 ml, 100 ml



COMPOSITION: 100 ml contains
 Florphenicol.....20 g
 Tylosin tartrate.....10 g
DOSAGE:
 I.M administration
 1 ml/ 20 kgs B.W.

INDICATION:
 For treatment of respiratory and digestive
 disease as pneumonia, APP, pasteurellosis,
 glasser, paratyphoid, *E.coli*.

COMPOSITION: 100 ml contains
 Spectinomycin HCl.....10 g
 Lincomycin HCl.....5 g
INDICATIONS:
 Against of mycoplasma, pneumonia,
 pasteurellosis, *E.coli*, *Salmonella*, diarrhea
 Treatment of mastitis, metritis, arthritis, hoof
 inflammation.

DOSAGE:
 For I.M, S.C administration
 - Cattle: 1 ml/ 10 kgs B.W.
 - Cats and dogs: 1 ml/ 5 kgs B.W.
 - Poultry: 1 ml/ 4 kgs B.W.

LINCO-S.1500

50 ml, 100 ml



MARCINE

50 ml, 100 ml



COMPOSITION: 100 ml contains
 Marbofloxacin.....5 g
INDICATION:
 Treatment of *Mycoplasma*, *Pasteurella*,
 atrophic rhinitis, CRD, CCRD, salmonellosis,
E.coli, arthritis, metritis.

DOSAGE:
 For I.M administration
 - Bovine: 1 ml/ 15 - 20 kgs B.W daily
 - Swine, goats, sheep: 1 ml/ 10 - 15 kg B.W
 daily.
 - Poultry: 1 ml/ 5 kgs B.W.

COMPOSITION: 100 ml contains
 1-cyclopropyl-7-(4-ethyl-1-piperazinyl)-6-flouro-
 1,4-dihydro-4-oxo-3-quinoline-carboxylic acid
 (enrofloxacin base).....10 g
INDICATIONS:
 Treat digestive diseases: diarrhea caused by
E.coli, bloody diarrhea and Salmonellosis,...

DOSAGE:
 - Adult cattle: 1 ml/ 15 - 20 kgs B.W.
 - Young cattle: 1 ml/ 10 - 15 kgs B.W.
 - Poultry: 1 ml/ 6 - 8 kgs B.W.

MAXLOX

50 ml, 100 ml





COMPOSITION: 100 ml contains
 Florphenicol.....20,000 mg
 Flunixin meglumine.....2,000 mg
INDICATION:
 For digestive and respiratory diseases, mastitis, metritis, secondary infection caused by PRRS, FMD.

DOSAGE:
 - Swine: 1 ml/ 16 - 22 kg B.W.
 - Buffalos, cattle, goats, sheep: 1 ml/ 19 - 30 kg B.W.
 - Poultry: 1 ml/ 10 - 12 kg B.W.

COMPOSITION: 100 ml contains
 Tiamulin HF.....10 g
 Colistin sulfate.....40,000,000 IU
INDICATIONS:
 For treatment of diarrhea, respiratory diseases, arthritis on livestock.

DOSAGE:
 I.M administration
 - Bovine, sheep: 1 ml/ 15 - 18 kg B.W for 3 days.
 - Swine: 1 ml/ 10 - 12 kg B.W for 3 days.



COMPOSITION: 100 ml contains
 Oxytetracycline HCl.....20 g
DOSAGE:
 For I.M or S.C administration
 General: 1 ml/ 10 kgs B.W

INDICATIONS:
 Arthritis, gastrointestinal and respiratory infections in calves, cattle, goats, sheep and swine.

COMPOSITIONS: 1 ml contains
 Norfloxacin.....50 mg
INDICATIONS:
 Prophylaxis and against of pneumonia, mastitis, metritis, pasteurellosis, diarrhea.

DOSAGE:
 - Swine, buffalos, cattle, goat, sheep: 1 ml/ 10 kgs B.W.
 - Poultry: 1ml/ 4 - 5 kgs B.W.



COMPOSITION: 100 ml contains
 Florfenicol.....10,000 mg
 Spiramycine adipate.....5,000 mg
 Prednisolone.....500 mg
 Lidocaine HCl.....1,000 mg
INDICATION:
 - Treatment of respiratory and digestive diseases in livestock.
 - Treatment of mycoplasma, C.CRD, edema, typhoid, pasteurellosis in poultry.

DOSAGE:
 I.M administration
 - Cattle: 1 ml/ 12 - 15 kgs B.W.
 - Swine, calves, goats, sheep: 1ml/ 8 - 10 kgs B.W.
 - Dogs, cats: 1ml/ 5 - 6 kgs B.W.
 - Poultry: 1 ml/ 5 - 6 kgs B.W.

COMPOSITION: 1 ml contains
 Oxytetracycline HCl.....300 mg
 Flunixin meglumine.....20 mg
INDICATION:
 Treatment of pneumonia reducing pyrexia in livestock.

DOSAGE:
 I.M or S.C administration
 1 ml/ 20 kgs B.W.



▶ Injectable solution

THIAM-LOSYN

50 ml, 100 ml



COMPOSITION: 100 ml contains
 Thiamphenicol.....20 g
 Tylosin tartrate.....10 g
INDICATIONS:
 For treatment of pneumonia, APP, pasteurellosis, paratyphoid, *E.coli*.

DOSAGE:
 I.M administration.
 - Higher 50 kg: 1 ml/ 20 kgs B.W.
 - Lower 50 kg: 1 ml/ 14 - 16 kgs B.W.
 - Dogs, cats, poultry: 1 ml/ 10 kgs B.W.

COMPOSITION: 100 ml contains
 Thiamphenicol.....10 g
 Oxytetracycline HCl.....5 g
 Dexamethasone sodium phosphate.....75 mg
 Bromhexin HCl.....100 mg
INDICATIONS: For treatment of digestive disease such as enteritis, diarrhea, paratyphoid, and respiratory diseases such as pasteurellosis, pneumonia. Against of metritis, mastitis, arthritis.

DOSAGE:
 I.M administration.
 - Bovine, swine: 1 ml/ 10 kgs B.W.
 - Piglets, calves, goats, sheep: 1 ml/ 7 - 8 kgs BW.
 - Dogs, cats, poultry: 1 ml/ 5 kgs B.W.

THIAM-SONE

50 ml, 100 ml



TILSYN

50 ml, 100 ml



COMPOSITION: 100 ml contains
 Tildipirosin.....18,000 mg
DOSAGE:
 S.C. administration
 1 ml/ 40 kgs B.W.
 Do not administer over 10 ml/ injection site.

INDICATION:
 Treatment and prevention of pneumonia caused by *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus sommi*.

COMPOSITION: 100 ml contains
 Tylosin tartrate.....20 g
DOSAGE:
 For I.M administration
 General dose: 1 ml/ 20 kgs B.W.

INDICATIONS:
 Against of *Campylobacter*, *Mycoplasma*, *Pasteurella*, *Staphylococcus*, *Streptococcus* and *Treponema* spp. in livestock.

TYLANVET

50 ml, 100 ml



TYLO-TETRA

50 ml, 100 ml



COMPOSITION: 1 ml contains
 Tylosin tartrate.....100 mg
 Oxytetracycline HCl.....100 mg
DOSAGE:
 I.M administration
 Sheep, goat, cattle: 1 - 1.5 ml/ 10 kgs B.W.

INDICATIONS:
 Treatment of metritis, mastitis, digestive infectious, pneumonia, shipping fever, secondary infection.

COMPOSITION: 100 ml contains
 Tylosin tartrate.....10 g
 Gentamycin sulfate.....5 g
INDICATIONS: Treatment of
 - Pneumonia, asthma caused by *Mycoplasma*, mastitis, metritis, ulcerated flesh...
 - CRD, sinusitis, pasteurellosis in poultry, livestock.
 - Arthritis in livestock & poultry, dysentery and other infected diseases.

DOSAGE:
 Once daily intramuscular injection for 3 - 4 days.
 - Bovine: 1 ml/ 20 - 30 kgs B.W.
 - Swine, goats, sheep: 1 ml/ 20 kgs B.W.
 - Dogs, cats, poultry: 1 ml/ 10 kgs B.W.

TYLO-GENTA

50 ml, 100 ml



AMGEN-S

50 ml, 100 ml



COMPOSITION: 100 ml contains
Amoxicillin trihydrate.....15,000 mg
Gentamycin sulfate.....4,000 mg
DOSAGE:
I.M administration
1 ml/ 10 kgs B.W.

INDICATION:
Enteritis, pneumonia, bronchitis, pleurisy, mastitis, metritis, nephritis, dermatitis, arthritis.

COMPOSITION: 100 ml contains
Ampicillin trihydrate.....10 g
Colistin sulfate.....40,000,000 IU
INDICATIONS:
Treatment of respiratory and digestive diseases.

DOSAGE:
S.C or I.M injection
- Calves, goats, sheep: 1 ml/ 10 kgs B.W.
- Bovine: 1 ml/ 15 kgs B.W.

AMPI-CS

50 ml, 100 ml



CEFOXYL

50 ml, 100 ml



COMPOSITION: 100 ml contains
Cefotaxime sodium.....10 g
INDICATIONS:
Treatment of meningitis, pneumonia, pasteurellosis, pleurisy, metritis, nephritis, arthritis.

DOSAGE: Deep intramuscular injection
- Piglet: 1 ml/ 10 kgs B.W.
- Pigs (>50 kg): 15 - 20 kgs B.W.
- Calves, goats, sheep: 1 ml/ 20 kg B.W.
- Bovines: 1 ml/ 30 kgs B.W.
- Poultry: 1 ml/ 8 kgs BW.

COMPOSITIONS: 1 ml contains
Ceftiofur HCl.....75 mg
Ketoprofen.....100 mg
INDICATIONS:
Treatment of pneumonia, respiratory disease, metritis.

DOSAGE:
I.M or S.C administration for 3 - 5 days.
- Dairy cow: 1 ml/ 25 - 35 kgs B.W.
- Cattle, buffalo, goat, sheep: 1 ml/ 40 - 50 kgs B.W.

CEPTIFEN

50 ml, 100 ml



CEQUIN 4

50 ml, 100 ml



COMPOSITION: 100ml contains
Cefquinome sulfate.....2.5 g
INDICATION:
Treatment of mastitis, metritis, pneumonia, asthma, digestive diseases, hoof inflammation.

DOSAGE:
I.M administration.
- Calves, foals: 1ml/ 12.5 kgs BW.
- Bovine, horses: 1 ml/ 25 kgs BW.

COMPOSITION: 100 ml contains
Ceftiofur HCl.....10 g
INDICATION:
Treatment of pleurisy, APP disease, mastitis, metritis, *E.coli*, paratyphoid.

DOSAGE:
I.M or S.C administration
- Piglet, dog, cat: 1 ml/ 15 - 20 kgs B.W.
- Small livestock: 1 ml/ 30 kgs B.W.
- Large livestock: 1 ml/ 50 kgs B.W.

CETIMAX

50 ml, 100 ml



▶ Injectable suspension

CLA-MOX

50 ml, 100 ml



COMPOSITION: 1 ml contains
Amoxicillin trihydrate.....140 mg
Clavulanic acid.....35 mg
INDICATIONS:
Treatment of respiratory, digestive and urinary disease.

DOSAGE:
I.M administration
1 ml/ 20 kgs B.W/ day.

COMPOSITION: 100 ml contains
Amoxicillin trihydrate.....10 g
Colistin sulfate.....40,000,000 IU
DOSAGE:
Daily I.M or SC administration
1 ml/ 10 kgs B.W for 3 days

INDICATIONS:
Treatment of diarrhea, enteritis erysipelas, M.M.A syndrome, paratyphoid, arthritis, respiratory tract infection.

COLIMOX

50 ml, 100 ml



NOVAMOX-LA

50 ml, 100 ml



COMPOSITION: 1 ml contains
Amoxiciline trihydrate.....150 mg
INDICATIONS:
Treatment of mastitis, metritis, abscess, arthritis, digestive and respiratory diseases.

DOSAGE:
I.M or S.C injection
1 ml/ 15 kgs B.W per day.

COMPOSITION: 1 ml contains
Dihydrostreptomycin sulfate.....200 mg
Benzathin penicillin G.....200,000 IU
INDICATIONS:
Treatment of arthritis, infectious case caused by Cocci, diarrhea, pasteurellosis, dermatitis, septicemia, abscess.

DOSAGE:
I.M administration
- Buffalos, cattle: 1 ml/ 20 - 30 kgs B.W.
- Goat, sheep, young livestock: 1 ml/ 15 - 20 kgs B.W.

PENISTREP

50 ml, 100 ml



TRIME-ZONE

50 ml, 100 ml



COMPOSITIONS: 100 ml contains
Sulfamethoxazole.....40 g
Trimethoprim.....8 g
INDICATIONS:
Treatment of typhoid, diarrhea, infectious diseases, pneumonia.

DOSAGE:
I.M administration.
General dose: 1 ml/ 25 - 30 kgs B.W/ day for 3 - 5 days

COMPOSITION: 100 ml contains
Ceftriaxone sodium.....10 g
INDICATION:
Treatment of cough, pleural-pneumonia, mastitis, metritis, agalactia.

DOSAGE: I.M administration
- Calves, goats, sheep, other young livestock: 1 ml/ 15 - 20 kgs/ B.W.
- Buffalos, cattle, other large livestock: 1 ml/ 30 kgs B.W.
- Poultry: 1 ml/ 8 kgs B.W.

TRIXON

50 ml, 100 ml





COMPOSITION: 10 ml contains
 Amoxicillin trihydrate.....200 mg
 Clavulanic acid.....50 mg
INDICATIONS:
 Treatment of mastitis in lactating period.

USAGE-INSTRUCTION:
 Should be infused into each affected quarter via the teat canal.

COMPOSITION: 10 ml contains
 Cephalexin monohydrate.....250 mg
 Neomycin sulfate.....250 mg
USAGE-INSTRUCTION:
 Should be infused into each affected quarter via the teat canal.

INDICATIONS:
 Treatment of clinical mastitis caused by sensitive microorganisms such as *Staphylococcus aureus*, *Streptococcus spp.*, *E. coli* in cows, goat and sheep during dry period.



COMPOSITION: 10 ml
 Cephalexin monohydrate.....200 mg
 Kanamycin monosulphate.....100,000 IU
USAGE-INSTRUCTION:
 Should be infused into each affected quarter via the teat canal.

INDICATIONS:
 Treatment and prevention of clinical mastitis caused by sensitive microorganisms such as *Staphylococcus aureus*, *Streptococcus spp.*, *E. coli* in cows, goat and sheep during lactating period.

COMPOSITIONS: 10 ml contains
 Ceftiofur hydrochloride.....500 mg
 Excipients q.s.....10 ml
USAGE-INSTRUCTION:
 Should be infused into each affected quarter via the teat canal.

INDICATIONS:
 Treatment of subclinical mastitis in dairy cattle, goats and sheep at dry period caused by sensitive microorganisms such as *Staphylococcus aureus*, *Streptococcus dysgalactiae*, *Streptococcus uberis* and *E.coli*.



COMPOSITION: 10 ml contains
 Ceftiofur hydrochloride.....125 mg
 Excipients q.s.....10 ml
USAGE-INSTRUCTION:
 Should be infused into each affected quarter via the teat canal.

INDICATIONS:
 Treatment of subclinical and clinical mastitis in dairy cattle, goats, sheep at lactating period caused by sensitive microorganisms such as *Staphylococcus aureus*, *Streptococcus dysgalactiae*, *Streptococcus uberis* and *E.coli*.



▶ Oral solution

CYLOSIN

500 ml, 1 liter



COMPOSITION: 100 ml contains
 Tylosin tartrate.....10 g
 Doxycyclin hyclate.....10 g
INDICATION:
 Treatment of asthma, diarrhea, sinusitis, typhoid, enteritis, diarrhea in livestock and poultry.

DOSAGE:
 - Poultry: 10 ml/ 10 - 12 liters of water or 10 ml/ 60 - 80 kgs B.W.
 - Livestock: 10 ml/ 80 - 100 kgs B.W.

COMPOSITION: 1 liter contains
 Florfenicol.....100 g
 Doxycyclin hyclate.....100 g
 Bromhexine HCl.....10 g
INDICATIONS:
 Treatment of *Actinobacillus* spp., *Pasteurella* spp., *Salmonella* spp., *Streptococcus* spp.

DOSAGE:
 - Poultry: 1 ml/ 1 liter of water or 1 ml/ 8 - 10 kgs B.W.
 - Livestock: 1 ml/ 1 - 2 liters of water or 1 ml/ 10 - 15 kgs B.W.

DOHEFLOR

500 ml, 1 liter



ENROHEXIN

500 ml, 1 liter



COMPOSITION: 100 ml contains
 Enrofloxacin base.....20 g
 Bromhexine HCl.....1.5 g
INDICATIONS:
 Against respiratory and digestive disease as gastrointestinal infection, diarrhea, CRD, C.CRD and Pasteurellosis.

DOSAGE:
 For 3 - 5 days.
 - Pigs, calves, goats & sheep: 1 ml/ 10 - 20 kgs B.W, twice per day
 - Poultry: 1 ml/ 3 - 4 liters water

COMPOSITION: 100 ml contains
 Enrofloxacin base.....10 g
INDICATION:
 Gastrointestinal, respiratory and urinary tract infections.

DOSAGE:
 - General dose: 1 ml/ 4 liters of water.
 - Calves, goats, sheep: 2 ml/ 15 - 30 kgs B.W.
 - Poultry: 2 ml/ 3 - 4 liters of water.
 - Other livestock: 2 ml/ 4 - 6 liters of water.

ENROVET

100 ml, 250 ml



FLOTYL

500 ml, 1 liter



COMPOSITION: 1 ml contains
 Tilmicosin phosphate.....100 mg
 Florfenicol.....100 mg
DOSAGE:
 1 ml/ 2 liters of water.

INDICATIONS:
 Against of *Actinobacillus* spp., *Pasteurella* spp., *Salmonella* spp., *Streptococcus* spp., etc.

COMPOSITION: 1 ml contains
 Tylosin tartrate.....50 mg
 Gentamycin sulfate.....25 mg
INDICATION:
 Treatment of CRD, pneumonia, pasteurellosis, digestive diseases.

DOSAGE:
 - Treatment
 - Poultry: 10 ml/ 6 - 8 liters of drinking water or 1 ml/ 4 - 5 kgs B.W.
 - Livestock: 1 ml/ 5 - 8 kgs B.W.
 - Prevention: ½ treatment dose

LOSINNET

500 ml, 1 liter





NEO-S

100 ml, 200 ml



COMPOSITION: 100 ml contains
 Neomycin sulfate.....6 g
 Colistin sulfate.....50,000,000 IU
INDICATION:
 Prevention and treatment of *E.coli*, gastro-
 enteritis, diarrhea, dysentery, paratyphoid.

DOSAGE:
 Shake well before use
 - Young animals: 1 ml/ 3 - 5 kgs B.W daily.
 - Mature animals: 1 ml/ 7 - 10 kgs B.W daily.

COMPOSITION: 1 ml contains
 Colistin sulfate.....200,000 IU
 Spectinomycin base.....50 mg
INDICATIONS:
 Treatment of *E.coli*, *Haemophilus*,
Mycoplasma and *Salmonella spp.*

DOSAGE:
 - Young livestock: 1 ml/3 - 5 kgs B.W.
 - Mature livestock: 1 ml/ 5 - 8 kgs B.W.

PEC-COLI

100 ml, 250 ml



PECTYL

100 ml, 250 ml



COMPOSITION: 100 ml contains
 Spectinomycin.....5 g
INDICATIONS:
 Treatment of enteritis disease.

DOSAGE:
 Oral administration
 1 - 2 ml/ head, after birth.

COMPOSITIONS: 100ml contains
 Enrofloxacin base.....2,500 mg
 Colistin sulphate.....20,000,000 IU
INDICATIONS:
 Treatment of gastrointestinal, respiratory
 and urinary tract infections.

DOSAGE:
 - Livestock: 1 ml/ 1 - 1.2 liters water or 1 ml/
 3 - 4 kgs B.W.
 - Poultry: 1 ml/ 0.5 liters water or 10 ml/ 20 -
 25 kgs B.W.

TREAVET

100 ml, 250 ml



TREXIM

500 ml, 1 liter



COMPOSITION: 100 ml contains
 Sulfadiazine 40 g
 Trimethoprim8 g
INDICATIONS:
 Treatment of digestive diseases as diarrhea,
 CRD, typhoid, enteritis, respiratory as
 pneumonia, pasteurellosis.

DOSAGE:
 - Poultry:
 + Prevention: 1 ml / 4 liters of water.
 + Treatment: 1 ml/ 2 liters of water or 1 ml/
 10 kgs B.W.
 - Livestock:
 + Prevention: 1 ml/ 20 kgs B.W.
 + Treatment: 1 ml/ 10 kgs B.W.

COMPOSITION: 100 ml contains
 Tilmicosin (as phosphate).....25 g
INDICATIONS:
 Treatment of *Mycoplasma spp.* *Pasteurella*
multocida, APP, *Actinomyces pyogenes* and
Mannheimia haemolytica.

DOSAGE:
 For oral administration
 - Poultry: 1 ml/3 liters of water.
 - Swine, calves: 1 ml/ 20 kgs B.W. or 8 ml/ 10
 liters of water.

VATIL-ORAL

500 ml, 1 liter



Powder for injection

STREPTONIN

16.3 g/ 100 ml



COMPOSITIONS: 1 bottle contains
 Penicillin G potassium.....10,000,000 IU
 Streptomycin sulfate.....10 g
INDICATION
 Against of mastitis, pneumonia, metritis,
 salmonellosis, arthritis, dermatitis,
 erysipelas suis.

DOSAGE:
 - Buffalos, cattle, horse: 1 ml/ 8 kgs B.W/ day
 - Swine, goats, sheep, calves: 1 ml/ 6 kgs B.W.
 - Poultry: 1 ml/ 4 kgs B.W.

COMPOSITION: 1 vial contains
 Ceftiofur sodium.....1 g
INDICATIONS:
 Prevention and treatment of diseases in poultry.

DOSAGE:
 - Fowl, duck:
 I.M or S.C administration: 1 ml/ 2 kgs B.W/ day.
 - Young ducks and chickens:
 SC administration: at 1 - 3 days of life: 1 ml/ 40 chicks.

TIOFURDIUM

1 g/ 100 ml



TRIXONE

10 g/ 100 ml



COMPOSITION: 1 vial contains
 Ceftriaxone sodium.....10 g
INDICATIONS:
 Treatment of infectious respiratory,
 pasteurellosis, typhoid, *E.coli*, etc.

DOSAGE:
 I.M injection
 - Piglet: 1 ml/ 10kgs B.W once daily.
 - Swine: 1 ml/ 20 kgs B.W once daily.
 - Poultry: 1ml/ 8 kgs B.W once daily.

Powder

COMPOSITION: 100 g contains
 Amoxicillin trihydrate.....50 g
DOSAGE:
 - Cattle: 2 g/ 100 kgs B.W, twice per day
 - Poultry: 200 g/ 1,000 liters of drinking water or 300 g/ 1 ton of feed.

INDICATIONS:
 - Livestock: Treatment of pneumonia, pleurisy, diarrhea
 - Poultry: Treatment of necrotic enteritis, diarrhea, umbilical infection, CRD.

AMOXY-WS

500 g, 1 kg



AUCOLIN

500 g, 1 kg



COMPOSITION: 100 g contains
 Amoxicillin trihydrate.....20 g
 Colistin sulfate.....120,000,000 IU
INDICATIONS:
 Treatment of diarrhea, pneumonia, arthritis, metritis, mastitis.

DOSAGE: For oral administration
 - Poultry:
 + Treatment: 1 g/ 2 liters of water or 100 g/ 1,000 kgs B.W.
 + Prevention: 1 g/ 4 liters of water.
 - Swine: 7 g/ 100 kgs B.W.
 - Cattle, buffalos: 20 g/ 400 kgs B.W.

COMPOSITION: 100g contains
 Streptomycin sulfate.....10,000 mg
 Berberin HCl.....2,000 mg
INDICATIONS:
 Treatment of enteritis, flatulence, indigestion, epidemic diarrhea, diarrhea.

DOSAGE:
 For oral administration
 - Cattle: 50 g/ 600 kgs B.W daily.
 - Poultry: 50 g/ 50 liters of water or 300 kgs B.W daily. Prevention: ½ dose of treatment for 2 - 5 consecutive days.

BERINTREP-WS

500 g, 1 kg



BIOCILIN CS

500 g, 1 kg



COMPOSITIONS: 100 g contains
 Ampicillin trihydrate..... 10 g
 Colistin sulfate.....40,000,000 IU
INDICATIONS:
 Treatment of respiratory, urinary tract infection in livestock and poultry.

DOSAGE:
 - Treatment:
 + Poultry: 1 g/ 1 liter of water or 10 g/ 50 kgs B.W.
 + Livestock: 300 g/ 200 kgs of feed.
 - Prevention: 1/2 treatment dose.

COMPOSITION: 100 g contains
 Cephalexin.....7.5 g
 Colistin sulfate.....20,000,000 IU
INDICATIONS:
 Against of *Staphylococcus*, *Streptococcus*, *Corynebacterium*, *Pasteurella*, *Salmonella*, *E.coli*, APP and *Mycoplasma* spp.,....

DOSAGE:
 - Cattle: 10 g/ 75 kgs B.W.
 - Poultry: 100 g/ 300 - 400 kgs B.W or 1 g/ 1 - 1.5 liters of water.

CEPLYTIN

500 g, 1 kg



COLIMAX-S

500 g, 1 kg



COMPOSITION: 100 g contains
 Colistin sulphate.....240,000,000 IU
INDICATIONS:
 Treatment of gastroenteritis, diarrhea, paratyphoid, necrotic enteritis in piglets; digestive infection.

DOSAGE:
 - Poultry: 1 g/ 4 liters of water or 100 g/ 2,000 kgs B.W.
 - Cattle: 10 g/ 400 kgs B.W daily.

COMPOSITION: 100 g contains
 Oxytetracyclin HCl.....6 g
 Colistin sulfate.....40,000,000 IU
INDICATIONS:
 Treatment for acute and chronic diarrhea, swollen head, enteritis.

DOSAGE:
 - Livestock: 10 g/ 70 - 80 kgs B.W.
 - Poultry: 10 g/ 30 - 40 kgs B.W.

COLIVIT.TETRA

500 g, 1 kg



COLYDOF

500 g, 1 kg



COMPOSITION: 100 g contains
 Doxycycline hyclate.....10 g
 Colistin sulfate.....120,000,000 IU
INDICATIONS:
 Against of *Mycoplasma hyopneumoniae*, *Pasteurella*, CRD, C.CRD, *Coryza*, diarrhea.

DOSAGE:
 - Calves, goats and sheep: Twice daily 1 g per 20 kg body weight.
 - Poultry: 1 kg per 2,000 liters of drinking water.
 - Swine: 1 kg per 1,000 liters of drinking water.

COMPOSITION: 100 g contains
 Lincomycin HCl.....10 g
 Colistin sulfate.....60,000,000 IU
DOSAGE:
 Oral administration
 Poultry: 1 g/ 2 liters of drinking water for 3 - 5 days

INDICATIONS:
 Against of *E. coli*, *Pseudomonas*, *Salmonella*, *Pasteurella*, *Staphylococcus*, *Streptococcus*, *Clostridium*, *Mycoplasma* spp.

COLYN-Z

500 g, 1 kg



Powder

COMPLEX AMIN

500 g, 1 kg



COMPOSITION: Each gram contains
 Erythromycin thiocyanate.....35 mg
 Oxytetracycline hydrochloride.....50 mg
 Streptomycin sulfate.....35 mg
 Colistin sulfate.....200,000 IU
 Vitamin A.....3,000 IU
 Vitamin D₃.....1,500 IU
 Vitamin E.....2 mg
 Vitamin B₁.....2 mg
 Vitamin B₂.....4 mg
 Vitamin B₆.....2 mg
 Vitamin B₁₂.....1 mcg
 Vitamin C.....20 mg

Ca - pantothenate.....10 mg
 Vitamin K₃.....2 mg
 Nicotinamide.....20 mg
 Inositol.....1 mg

INDICATIONS:

- Stimulation of egg production, increasing growth, improving feed conversion and vitamin supplement.
 - Treatment of gastrointestinal, respiratory and urinary infections

DOSAGE:

For oral administration through feed.
 Poultry and swine: 100 g/ 100 liters of water or 100 g/ 50 kgs of feed.

COMPOSITION: 100 g contains

Tilmicosin phosphate.....20,000 mg
 Doxycycline hyclate.....20,000 mg

INDICATIONS:

Control and treatment of respiratory infections.

DOSAGE:

For oral administration.
 - Poultry: 1 g/ 3 - 4 liters of water.
 - Swine: 2 g/ 100 kgs B.W or 265 g/ 1 ton of feed.

CYCOSIN

500 g, 1 kg



CYN-DOX

500 g, 1 kg



COMPOSITION: 100 g contains

Doxycycline hyclate.....10 g
 Neomycin sulfate.....10 g

DOSAGE:

- Livestock: 10 g/ 100 kgs B.W/ day
 - Poultry: 10 g/ 50 kgs B.W/ day or 1 g/ 1 - 2 liters of drinking water.

INDICATIONS:

For treatment of *Mycoplasma hyopneumoniae*, *Pasteurella*, *E.coli*, *Salmonella*, *Staphylococcus*, *Streptococcus*, CRD, C.CRD, Coryza, ...

COMPOSITION: 100 g contains

Tylosin tartrate.....10 g
 Doxycycline hyclate.....10 g

INDICATIONS:

Against of *Bordetella*, *Campylobacter*, *Chlamydia*, *E.coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus*, *Streptococcus* and *Treponema*,...

DOSAGE:

For oral administration
 - Swine, poultry: 20 g/ 200 kgs B.W per day, equivalent 20 g/ 20 - 40 liters of water
 - Calves, kids, lambs: twice daily 20 g/ 200 kgs B.W.

DOXLOSIN

500 g, 1 kg



DOXYFLOR

500 g, 1 kg



COMPOSITION: 100 g contains

Florfenicol.....10,000 mg
 Doxycycline hyclate.....10,000 mg
 Bromhexine HCl.....1,000 mg

INDICATIONS:

Against of pneumonia, salmonellosis, pasteurellosis, fever, diarrhea, enteritis, metritis, mastitis, swollen head.

DOSAGE:

- Prevention:
 + Poultry, piglets: 1 g/ 4 liters of water or 1 g/ 2 kgs of feed or 1 g/ 30 - 40 kgs B.W.
 + Pigs, calves: 1 g/ 6 liters of water or 1 g/ 3 kgs of feed or 1 g/ 60 - 80 kgs B.W.
 - Treatment: double dose of prevention.

COMPOSITION: 100 g contains

Doxycycline hyclate.....50 g

DOSAGE:

For oral administration
 - Bovine: Twice daily 5 g/ 200 kgs B.W.
 - Poultry and swine: 1 g/ 2 - 4 liters of water.

INDICATIONS:

Treatment of respiratory and digestive tract infection caused by *Bordetella*, *Campylobacter*, *Chlamydia*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella*, *Rickettsia*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. on livestock.

DOXYVET

500 g, 1 kg



ERYCIN-B

500 g, 1 kg



COMPOSITION: 100 g contains
 Erythromycin.....5,000 mg
 Sulfamethoxyipyridazine.....10,000 mg
INDICATION:
 Treatment of pneumonia, pasteurellosis, CRD, enteritis, ...

DOSAGE:
 - Treatment:
 + Poultry: 2 g/ 1 liter of water or 1 g/ 5 kgs B.W/ day
 + Livestock: 1 g/ 10 kgs B.W/ day or 100 g/ 50 - 100 kgs of feed
 - Prevention: ½ treatment dose

COMPOSITION: 100 g contains
 Amoxicillin trihydrate.....10 g
 Gentamycin sulfate.....3 g
INDICATIONS:
 Treatment of respiratory disease, diarrhea, salmonellosis, pasteurellosis.

DOSAGE:
 - Poultry: 10 g/ 8 - 10 liters of water or 10 g/ 50 kgs B.W.
 - Livestock: 10 g/ 100 kgs B.W.

GENTACIN

500 g, 1 kg



LEXYN

500 g, 1 kg



COMPOSITION: 100 g contains
 Cephalixin.....20 g
INDICATIONS:
 Treatment of *Staphylococcus*, *Streptococcus*, *Corynebacterium*, *Pasteurella*, *Salmonella*, *E.coli*, *Actinobacillus* spp.,...

DOSAGE:
 - Cattle: 10 g/ 200 kgs B.W.
 - Poultry: 10 g/ 1000 kgs B.W or 1 g/ 4 liters of drinking water.

COMPOSITION: 100 g contains
 Enrofloxacin HCl.....20 g
DOSAGE:
 For oral administration
 - Bovine: 1 g/ 60 kgs B.W for 3 - 5 days.
 - Swine: 1 g/ 40 kgs B.W for 3 - 5 days.
 - Poultry: 1 g/ 4 liters of water for 3 - 5 days.

INDICATIONS:
 - Against of pasteurellosis, *Mycoplasma*, diarrhea, salmonellosis, enteritis, ..in livestock.
 - Against of pasteurellosis, coryza, CRD, C.CRD, diarrhea, salmonellosis, dysentery, enteritis in poultry.

MAXFOC-WS

500 g, 1 kg



MOXCYCIN

500 g, 1 kg



COMPOSITION: 100 g contains
 Amoxicillin trihydrate.....14 g
 Bromhexine HCl.....0.5 g
INDICATIONS:
 Effectively treat *E.coli*, salmonellosis, enzootic pneumonia, pasteurellosis.

DOSAGE:
 - Poultry: 1 g/ 1.5 liters of water or 1 g/ 8 - 10 kgs B.W.
 - Swine: 1 g/ 15 kgs B.W.
 - Cattle: 1 g/ 30 kgs B.W.

COMPOSITION: 1 g contains
 Neomycin sulphate.....200 mg
INDICATION:
 Against of enteritis.

DOSAGE:
 - Calve, goat, sheep: 50 mg/ kg B.W.
 - Poultry and swine: 300 g/ 500 - 600 liters of drinking water.

NEOCIN-WS

500 g, 1 kg



 Powder

NEOTYN

500 g, 1 kg

**COMPOSITION:** 100 g contains

Neomycin sulfate.....6 g
Colistin sulfate.....36,000,000 IU

INDICATIONS:

- Swine: gastroenteritis, salmonellosis, diarrhea, dysentery, swollen head.
- Poultry: omphalitis, diarrhea, salmonellosis, pasteurellosis.

DOSAGE:

- Poultry: 1 g/ 2 - 3 kgs B.W or 2 g/ 1 liter of water for 3 days.
- Swine: 10 g/ 30 - 40 kg B.W, twice per day
- Piglet under a month of age: 10 g/ 3 piglets, twice per day, for 3 days.

COMPOSITION: 100 g contains

Oxytetracycline HCl.....18 g
Neomycin sulfate.....10 g

INDICATIONS:

Against of respiratory infections, CRD, typhoid, gastrointestinal infections.

DOSAGE:

For mixing with feed or water for 3 - 5 days.

- Treatment:

+ Cattle: 1 g/ 10 - 12 kgs B.W/ day.
+ Pigs: 1 g/ 8 - 10 kgs B.W/ day.
+ Poultry: 1g/ 10 - 12 kgs BW/ day or 1 g/ 1 liter of water/ day.
+ Bees: 1 g/ 0.5 liter of syrup/ bees.
- Prevention: ½ treatment dose.

NEOX

500 g, 1 kg

**COMPOSITION:** 100 g contains

Oxytetracycline HCl.....20,000 mg
Vitamin A.....100,000 IU
Vitamin D₃.....20,000 IU
Vitamin E.....30 IU
Vitamin B₁₂.....400 mcg
Vitamin K.....40 mg
Riboflavin.....160 mg
Niacinamid.....800 mg
Pantothenic acid.....300 mg

INDICATIONS:

- Enteritis, pasteurellosis, synovitis, sinusitis...
- Improve digestion, increasing growth rate.

DOSAGE:

For oral administration

- Prevention: 1 g/ 4 liters of water, equivalent 0,15 g/ 4 kgs B.W for 3 days.
- Treatment: 2 g/ 4 liters of water, equivalent 0,3 g/ 1 kg B.W for 5 days.

NITAXY

500 g, 1 kg

**COMPOSITION:** 100 g contains

Oxolinic acid.....25,000 mg

INDICATIONS:

C.CRD, enteritis, typhoid, pasteurellosis, swollen head.

DOSAGE:

For oral administration

- Prevention: 1 g/ 2 liters of drinking water or 7.5 g/ 150 kgs B.W/ day.
- Treatment: 1 g/ 1 liter of drinking water or 15 g/ 100 kgs B.W/ day.

NIXOLIN

500 g, 1 kg

**COMPOSITION:** 100 g contains

Oxytetracycline HCl.....5,500 mg
Vitamin A.....660,000 IU
Vitamin D₃.....117,000 IU
Vitamin E.....660 IU
Vitamin B₁₂.....1,100 mcg
Vitamin K.....440 mg
Riboflavin.....216 mg
Vitamin PP.....2,760 mg
Pantothenic acid.....810 mg

Pyridoxine.....36 mg
Sodium Chloride.....4,250 mg
Potassium Chloride.....1,700 mg

INDICATIONS: Increase eggs quantity, productivity and hatch rates. Decrease mortality. Improve egg-shell quality. Protect animals from diseases, maintain eggs quantity.

DOSAGE:

- 1g/ 4 liters of water, used daily.
- 1 g/ 3.2 liters of water during laying period.

OXIMIN

500 g, 1 kg

**COMPOSITION:** 100 g contains

Spiramycin adipate.....9,600,000 IU
Oxytetracyclin HCl.....10 g

INDICATIONS:

Treatment of CRD, digestive diseasea.

DOSAGE:

Treatment: 2 g/ 1liter drinking water

OXPATE

500 g, 1 kg





COMPOSITIONS: 1 kg contains
 Lincomycin HCl.....200 g
 Spectinomycin HCl.....200 g
INDICATION:
 Treatment and prevention of digestive infection and respiratory infection.

DOSAGE:
 - Treatment
 + Poultry: 1 g/ 3 - 4 liters of drinking water or 1 g/ 15 - 20 kgs B.W.
 + Livestock: 1 g/ 5 liters of drinking water or 1 g/ 30 - 40 kgs B.W.
 - Prevention: ½ treatment dose.

COMPOSITION: 100 g contains
 Josamycin.....900,000 IU
 Oxytetracycline HCl.....15 g
INDICATIONS:
 Against of pneumonia, respiratory diseases, stimulate immunological system after vaccination.

DOSAGE:
 Treatment:
 - Livestock: 100 g/ 50 kgs of feed or 100 g/ 1,500 kgs B.W.
 - Poultry: 100 g/ 25 kg of feed or 100 g/ 750 kgs of B.W.
 Prevention: 1/2 treatment dose.



COMPOSITION: 100 g contains
 Josamycin.....18,000,000 IU
 Trimethoprim.....10 g
INDICATION:
 Treatment of chronic respiratory diseases, *E.coli*, *Salmonella*.

DOSAGE:
 - Treatment: 2 g/ 5 liters of water or 100 g/ 200 kgs B.W.
 - Prevention: 1/2 treatment dose.

COMPOSITION: 1 kg contains
 Josamycin.....18,000,000 IU
INDICATION:
 Against of pneumonia, stimulate of immunological system after vaccination.

DOSAGE:
 - Treatment: 2 kg/ ton of feed or 100 g/ 1,500 kgs B.W.
 - Prevention: 1/2 treatment dose.



COMPOSITION: 100 g contains
 Ampicillin trihydrate.....7.5 g
 Sulfamethoxazole.....12 g
INDICATIONS:
 Treatments of diarrhea, paratyphoid, necrotic enteritis.

DOSAGE:
 For 3 - 4 consecutive days
 - Poultry: 1 g/ 1 liters of water or 10 g/ 20 kgs of feed.
 - Swine: 1 g/ 10 kgs B.W or 100 g/ 40 kgs of feed.

COMPOSITION: 100 g contains
 Oxytetracycline HCl.....10,000 mg
 Paracetamol.....10,000 mg
 Vitamin C.....10,000 mg
INDICATIONS:
 Against of influenza, pain relief, reducing fever and infectious diseases.

DOSAGE:
 - Cattle: 1 g/ 5 kgs B.W.
 - Poultry: 1 g/ 1 liter of water.



Powder

TETROZ

500 g, 1 kg



COMPOSITION: 1 g contains
Oxytetracyclin HCl.....100 mg
INDICATIONS:
Against of *Bordetella*, *Bacillus*, *E. coli*,
Pasteurella, *Salmonella*, *Staphylococcus* and
Streptococcus spp.,....

DOSAGE:
For oral administration
- Livestock: 10 g/ 80 - 100 kgs B.W.
- Poultry: 1 g/ 2 - 3 liters of water.

COMPOSITION: 100 g contains
Sulfamethoxazol.....40 g
Trimethoprim.....8 g
INDICATIONS:
Against of *E. coli*, *Haemophilus*, *Pasteurella*,
Salmonella, *Staphylococcus* and
Streptococcus spp.

DOSAGE:
For oral administration.
- Calves, goats and sheep: Twice daily 5 g/
100 kgs B.W.
- Poultry and other livestock: 2 kg/ 3,000 -
5,000 liters of drinking water or 1 g/ 10 -
12 kgs B.W.

THOXAPRY

500 g, 1 kg



TIMIVET

500 g, 1 kg



COMPOSITION: 100 g contains
Tilmicosin phosphate.....20 g
INDICATIONS:
Against of respiratory infections

DOSAGE:
- Prevention:
+ Poultry: 1 g/ 4 - 5 liters of water or 1 g/ 20 -
30 kgs B.W.
+ Livestock: 1 g/ 40 - 50 kgs B.W or 1 kg/ 1 -
1.5 tons of feed
- Treatment: Double of prevention dose.

COMPOSITIONS: 100 g contains
Bacitracin methylene disalicylate.....50 g
INDICATIONS:
Against of necrosis enteritis.

DOSAGE:
- Poultry: 1 g/ 7 - 10 liters of water.
- Swine: 100 g/ 2,500 - 3,000 kgs B.W.

TRACINNET

500 g, 1 kg



TYCOCIN

500 g, 1 kg



COMPOSITIONS: 1 kg contains
Spiramycin adipate.....6,400,000 IU
Tylosin tartrate.....100 g
Colistin sulfate.....55,000,000 IU

INDICATIONS:
Against of colibacillosis, sinusitis, CRD,
synovitis and other respiratory infections.
DOSAGE:
100 g/ 150 - 200 liters of drinking water or
100 g/ 60 kgs of feed.

COMPOSITION: 100 g contains
Tylosin tartrate.....14,000 mg
Gentamycine sulfate.....7,000 mg
INDICATIONS:
- Against of *Mycoplasma*, pleurisy, pneumonia,
pasteurellosis in livestock.
- Against of CRD, C.CRD, diarrhea, necrosis
enteritis, respiratory disease, swollen face in poultry.

DOSAGE:
- Treatment:
+ Poultry: 10 g/ 14 liters of water of
100 g/ 30 - 40 kgs of feed.
+ Cattle: 10 g/ 200 kgs B.W of 100 g/
60 - 100 kgs of feed.
- Prevention: ½ dose of treatment.

TYLAGEN

500 g, 1 kg





COMPOSITION: 100 g contains
Tylosin (tartrate).....50 g

INDICATIONS:
- Poultry: CRD, sinusitis...
- Livestock: pneumonia,
asthma, dysentery, arthritis,
hoof inflammation.

DOSAGE:

- Poultry:
+ Prevention: 100 g/ 200 liters of water.
+ Treatment: 100 g/ 100 liters of water or 1 g/ 20 kgs B.W.
- Livestock:
+ Prevention: 100 g/ 200 liters of water or 100 g/ 100 kg
of feed.
+ Treatment: 100 g/ 100 liters of water or 100 g/ 50 kgs of
feed or 1 g/ 50 kgs B.W.

COMPOSITION: 100 g contains
Oxytetracyclin HCl.....20 g
Tylosin tartrate.....4 g

INDICATIONS:
Treatment of respiratory disease and
digestive disease: enteritis, diarrhea.
Arthritis, dermatitis, mastitis.

DOSAGE:

For oral administration
- Prevention: Daily 2.5 g/ 20 kgs B.W.
- Treatment: Daily 5 g/ 20 kgs B.W



COMPOSITION: 100 g contains
Streptomycin sulfate..... 5,000 mg
Neomycin sulfate..... 5,000 mg
Vitamin B₁.....300 mg
Vitamin B₂.....3 mg
Vitamin B₅.....200 mg
Vitamin B₆.....200 mg
Vitamin B₁₂..... 1,000 mcg
Vitamin PP..... 100 mg

INDICATIONS:
Prevent and treat diarrhea.

DOSAGE:
- Under 15 days of age: twice daily 1 - 1.5 g.
- Over 15 days of age: twice daily 2 g.

COMPOSITION: 100 g contains
Levofloxacin HCl.....10 g
Colistin sulfate.....10 g

DOSAGE:
- Treatment: 1 g/ 2 liters of water.
- Prevention: ½ treatment dose.

INDICATION:
Treatment and prevention of pneumonia,
sinusitis, bronchitis, mastitis, enteritis,
pericarditis, CRD, CCRD, coryza.



Injectable solution



COMPOSITION: 100 ml contains
Ivermectin.....1,000 mg

INDICATIONS:
Treatment of gastrointestinal roundworms and lungworm infections, lice,...

DOSAGE:
For S.C administration
- Bovine, horse: 1 ml/ 40 - 50 kgs B.W.
- Goats, sheep, other livestock: 1 ml/ 20 - 30 kgs B.W.
- Prevention dose: once every three months.

COMPOSITION: 1 ml contains
Doramectin.....10 mg

INDICATIONS:
Treatment of roundworm on digestive tract, lungworm, eyeworm, scabies, mites, lice, maggots.

DOSAGE:
- Dogs, cats: S.C administration
+ Internal parasites treatment: 1 ml/ 10 - 15 kgs B.W.
+ External parasite treatment: 1 ml/ 25 - 30 kg BW
- Livestock: 1 ml/ 50 kgs B.W for S.C or I.M administration



COMPOSITION: 100 ml contains
Ivermectin.....500 mg
Nitroxylnil.....25 g

DOSAGE: S.C administration
1 ml/ 25 kg B.W.

INDICATION:
Prevention and treatment of internal parasites: roundworm, liver fluke, lungworm, eyeworm.

COMPOSITION: 1 ml contains
Closantel sodium.....100 mg
Ivermectin.....8 mg

INDICATIONS:
Treatment of internal parasitosis and external parasitosis (lice, scabies, flies, among others).

DOSAGE:
S.C. administration
- Cattle: 1 ml/ 40 kgs of B.W.
- Sheep, camel and goats: 1 ml/ 20 kgs of B.W.



COMPOSITIONS: 1 ml contains
Ivermectin.....10 mg
Clorsulon.....100 mg

DOSAGE: S.C administration
1 ml/ 50 kgs B.W.

INDICATIONS:
Treatment of internal and external parasites: roundworm, worm on lung, eye worm, maggot, scabies, lice.

Oral solution

COMPOSITION: 100 ml contains
Diclazuril.....1 g

INDICATIONS:
Coccidiosis by killing some parasites.

DOSAGE:
Poultry: 1 ml/ 10 kgs BW or 500 - 800 ml/ 1,000 liters of water





COCUS TREAT

500 ml, 1 liter



COMPOSITION: 100 ml contains
 Toltrazuril.....2,500 mg
DOSAGE:
 1 ml/ 1 liter of water

INDICATIONS:
 Treatment and prevention of coccidiosis on livestock and poultry.

COSISTRE

500 ml, 1 liter



COMPOSITION: 100 ml contains
 Fluconazole.....1,500 mg
INDICATIONS:
 Treatment mycosis on dogs, cats and poultry.

DOSAGE:
 - Treatment: 1 ml/ 5 kgs B.W or 1ml/ 1 liter of water.
 - Dogs, cats: 1 ml/ 4 - 6 kgs B.W, 1 time/ day.

DAZONE

500 ml, 1 liter



COMPOSITION: Each ml contains
 Albendazole.....100 mg
INDICATIONS:
 Deworming, antihelminth.

DOSAGE: For oral administration
 - Swine, calves, cattle, buffalos: 1 ml/ 12 kgs B.W.
 - Sheep, goat: 1 ml/ 20 kgs B.W.
 - Poultry: 1ml/ 8 - 10 kgs B.W or 1 ml/ 1 liter of water.
 - Dogs, cats: 1 ml/ 5 - 10 kgs B.W.

DEWOM

500 ml, 1 liter



COMPOSITION: 100 ml contains
 Fenbendazole.....10g
INDICATION:
 Prophylaxis and treatment of gastrointestinal and respiratory worm infections.

DOSAGE:
 - Goats, swine and sheep: 1 ml per 20 kgs body weight.
 - Calves and cattle: 7.5 ml per 100 kgs body weight.
 - Poultry: 1 ml/ 4 - 6 kgs B.W.

SUPERIL

500 ml, 1 liter



COMPOSITION: 100 ml contains
 Toltrazuril.....5g
INDICATIONS:
 Prevention and treatment of coccidiosis.

DOSAGE:
 - Piglets at 3rd day after birthing: only dose, 0.4 ml/ 1 kg B.W or 1 ml/ piglet
 - Calves at the 5th week after birthing: only dose, 3 ml/ 10 kgs B.W.

WOR-F

500 ml, 1 liter



COMPOSITIONS: 100 ml contains
 Ivermectin.....1,000 mg
INDICATIONS:
 Treatment of external parasites.

DOSAGE:
 Poultry: Not more than 1 ml/ 10 kgs B.W, pour along spine.

▶ External solution

VEROTIN

100 ml, 250 ml



COMPOSITIONS: 1 ml contains
Ivermectin.....5 mg

INDICATIONS:
Treatment of diseases caused by internal and external parasites.

DOSAGE:
For topical administration
Cattle: 1 ml/ 10 kg B.W.

COMPOSITION: 1 ml contains
Phoxim.....75 mg

INDICATIONS:
Against of mange, mites, flea, lice, external parasites.

DOSAGE:
* Tapping medicine from dorsal to tail.
- Calve, goats, sheep, young livestock:
+ 4 ml/ animal (under 10 kgs B.W)
+10 - 20 ml/ animal (over 10 kgs B.W)
- Large livestock, cattle, buffalos: 20 - 30 ml/ animal
* Dilute into water, spraying: 1 ml/ 100 - 150 ml

XIMO

100 ml, 250 ml



▶ Powder

ADE SULTRY

500 g, 1 kg



COMPOSITIONS: 100 g contains
Sulfachloropyrazine.....7,500 mg
Trimethoprim.....1,500 mg
Vitamin A.....100,000 IU
Vitamin D₃.....10,000 IU
Vitamin E.....10 IU

INDICATION:
Protozoal infections specially coccidiosis in poultry. Gastrointestinal hemorrhage.

DOSAGE:
1 g/ 1 liter of drinking water

COMPOSITION: 100 g contains
Sulfachloropyridazine sodium.....30 g

DOSAGE:
100 g/ 120 - 150 kgs of feed or 1 g/ 2 liters of drinking water.

INDICATIONS:
Against of intestinal coccidiosis, diarrhea, and salmonellosis.

COX-STOP

500 g, 1 kg



IVER-WS

500 g, 1 kg



COMPOSITION: 1 g contains
Ivermectin.....3 mg

DOSAGE:
- Swine: 1 g/ 30 kgs B.W for 7 days.
- Calves, goats, sheep: 1 g/ 15 kgs B.W. for 2 - 3 days.

INDICATION:
Treatment of internal parasites as roundworms, pinworms, lung worms, trichocephalus, strongyloidiasis..., parasites in the intestines, kidneys, lungs, and external parasites as ticks, lice, scabies.

COMPOSITION: 100 g contains
Albendazol.....10 g
Levamisol HCl.....2 g

INDICATIONS:
Kills parasitic, worms in digestive and respiratory systems.

DOSAGE:
- Poultry: 10 g/ 100 kgs B.W.
- Small livestock: 10 g/ 150 kgs B.W.
- Large livestock: 10 g/ 400 kgs B.W.

LEVAZOL

500 g, 1 kg





NYTANZ

500 g, 1 kg



COMPOSITIONS: Each 100 g contains
Nystatin.....10,000,000 IU
INDICATION:
Treatment of infections caused by fungus.

DOSAGE: Mix with feed or water
- Poultry: 100 g/ 200 - 300 kgs B.W/ day
- Swine, calves, sheep, goats: 100 g/ 300 - 400 kgs B.W/ day.
- Bovine: 100 g/ 1,000 kgs B.W/ day.

COMPOSITION: 1 g contains
Amprolium hydrochloride.....300 mg
INDICATIONS:
AMPROLIN is indicated for coccidiosis.

DOSAGE: For oral administration
- Calves, sheep and goats:
+ Preventive: 1 g/ 60 kgs B.W.
+ Curative: 1 g/ 30 kgs B.W.
- Poultry:
+ Preventive: 1 kg/ 5,000 liters of drinking water.
+ Curative: 1 kg/ 1,250 - 2,500 liters of drinking water.

PROLICOX

500 g, 1 kg



PROXALIN.K

500 g, 1 kg



COMPOSITIONS: 1 kg contains
Amprolium HCl.....200 g
Sulfaquinoxaline.....150 g
Vitamin K₃.....5 g
INDICATIONS:
Treatment of coccidiosis in livestock and poultry.

DOSAGE: Oral administration
* Treatment dose for 5 - 7 days
- Poultry: 1 g/ 1,5 - 2 liters of drinking water or 1 g/ 8 - 10 kgs B.W.
- Calves, sheep and goats: 1 g/ 8 - 12 kgs B.W.
* Prevention dose for 21 days
10 g/ 50 - 100 kgs feed or 10 g/ 500 - 800 kgs B.W.

COMPOSITION: 100 g contains
Levamisol HCl.....10,000 mg
INDICATIONS:
Deworming in livestock and poultry.

DOSAGE:
- Livestock: 2 g/15kgs B.W.
- Poultry: 1 g/ 5 kgs B.W.

VAMISONE

500 g, 1 kg



VERYDINE

500 g, 1 kg



COMPOSITION: 100 g contains
Sulfadimidine sodium.....20 g
Diaveridine HCl.....2 g
INDICATIONS:
Prevention and treatment of cecal and intestinal coccidiosis, diarrhea, salmonellosis in animals.

DOSAGE:
For oral administration.
- Treatment: 1 g/ 1 litre of water for 3 - 5 consecutive days. Dosage intervals 2 days.
- Prevention: 1 g/ 1.5 litre of water for 3 - 5 consecutive days.

COMPOSITION: 100 g contains
Fenbendazole.....2,000 mg
MgSO₄.....2,000 mg
INDICATIONS:
Anthelmintic, deworming on digestive tract, lung,..

DOSAGE:
In chickens:
- Single dose of treatment: 25 g/ 50 kgs B.W.
- The frequent treatment: 25 g/ 2.5 kgs B.W.
In cattle: The only treatment: 38 g/ 100 kgs B.W.

ZOBENVET

500 g, 1 kg



ANTIBACTERIAL SPRAY

200 ml



COMPOSITION: 200 ml contains Oxytetracycline HCl.....5 g
INDICATIONS:
 Specific treatments for hoof inflammation, sole ulcers, injuries caused by castration and scabies.

DOSAGE:
 - Clean wounds and then spray on wounds, ulcers.
 - Use the medicine twice daily for 3 - 5 consecutive days

COMPOSITION: 1 liter contains Glutaraldehyde.....75,000 mg Alkyl Dimethyl Benzyl Ammonium Chloride.....150,000 mg
INDICATIONS:
 DYTAN is a broad-spectrum disinfectant to viruses, bacteria, spores, *Mycoplasma* and fungi causing disease on.

DOSAGE:
 - Disinfection of cages: 25 ml/ 10 liters of water.
 - Disease outbreaks: 33 ml/ 10 liters of water
 - Disinfection in facilities, incubators: 20 ml/ 10 liters of water.
 - Disinfection of carcass, animal fertilizers, boot dips: 100 ml/ 10 liters of water.
 - Disinfection of eggs: 10 ml/ 10 liters of water.
 - Disinfection of water: 5 ml/ 10 liters of water.

DYTAN

1 liter, 5 liters



OXIDAN

500 g, 1 kg



COMPOSITIONS: 100g contains Peroxygen (Potassium monopersulfate triple salt).....49.8% Organic acid (Malic acid).....9.3% Inorganic buffer system (Sodium hexametaphosphate, Sodium chloride, Sulfamic acid); Surfactant (Sodium dodecyl benzene sulphonate) q.s.....100 g
INDICATIONS: Disinfection of facilities, equipment, vehicles, disinfectant boot bath, water pipeline system, all surface in the farm. Disinfection of hard surface in livestock production and transportation facilities. Treatment of environment which contaminate organic compound.

DOSAGE:
 - Disinfectant of water system: Mix 100 g OXIDAN in 20 liters water (1:200), fill the solution into water system, keep into 1 hour, rinse the water system with water.
 - Sterilize drinking water and skin antiseptic: Mix 100 g OXIDAN in 100 liters of water (1:1,000) for drinking or bathing.
 - Air disinfection: when the disease are spreading, mix 100 g in 20 liters of water (1: 200), spray directly on animals daily with 1 liter/ 10 m² surface to prevent disease spreading.
 - Sterilize equipment: Mix 100 g of water in 10 - 20 liters of water (1: 100 - 1: 200), spray 300 ml/ m².
 - Sterilize housing before animal introduction: Mix 100 g in 10 liters of water (1: 100), spray 300 ml/ m².
 - Dip shoes, wheels: Mix 100 g in 10 liters of water (1:100).


COMPOSITION: 100 ml contains Povidone Iodine.....10 g
INDICATIONS:
 Disinfectant cage, vehicle, housing.

DOSAGE:
 - Disinfection of cages, equipment, vehicles: 10 ml/ 4 liters of water.
 - Disinfection of breast, uterus: 10ml/ 5 liters of water.
 - Wound: 10 ml/ 2.5 liters of water.
 - Hatchery disinfection: 10 ml/ 5 liters of water.

VODINE

1 liter, 5 liters





Quality affirms brand

Quality affirms brand



AIZ International Co.,LTD

 93-1034G Le Duc Tho Str., Ward 6, Go Vap Dist., HCMC, Vietnam

 (+84.274) 377 6891

 (+84.274) 377 6895

 sales@aizinternational.com

 www.aizinternational.com