


AMOXYCLAV



COMPOSITION

Each 1 ml contains

Amoxicillin (as trihydrate)	140mg
Clavulanic acid	35mg

INDICATIONS

For treatment of animal bacterial infection which is sensitive to Amoxicillin, such as pasteurellosis, pneumonia, mastitis, metritis, pullorum, salmonellosis, sepsis and so on.

ADMINISTRATION AND DOSAGE

In cattle, swine dogs and cats: Administer intra-muscularly or subcutaneously at a dosage of 1ml per 20kg of body weight.

Once a day, use for 3 – 5 days.

ADVERSE REACTIONS

A small number of animals may cause pain or local tissue reactions after injected.

PRECAUTIONS

Shake well before using and gently massage the injection site after injected. People with hypersensitivities (allergies) to penicillins should not handle the product.

If occur allergic reactions just from contact, such as rash, severe symptoms of the face, eyes or lips swelling and difficult breathing, please contact your doctor immediately. Avoid contact with the drug when operating. Use completely dry needles and syringes to prevent the water droplets contaminated remaining injection. Please used in 28 days after opening.

Discard expired drugs in accordance with the provisions of the local environmental protection departments. Keep out of reach of children.

WITHDRAWAL TIME

Swine and cattle: 14 days.
Milk: 60 hours

STORAGE AND EXPIRED TIME

Store in below 30°C, and protect from light.
3 years.

PACKING

50ml, 100ml

FLUN-50



COMPOSITION

Each 1 ml contains
Flunixin Meglumine

50mg

INDICATIONS

It is used for febrile and inflammatory diseases of livestock and small animals, muscle pain and soft tissue pain, such as sow-free syndrome in sows.

DOSAGE AND ADMINISTRATION

For intramuscular or intravenous injection.
One dose, cattle and swine: 1ml per 25kg bodyweight
dog and cat: 0.2ml- 0.4ml per 10kg bodyweight, once or twice a day for not more than 5 days.

PRECAUTION

Use with caution in animals with a history of gastrointestinal ulcers, kidney disease, liver disease or blood disease
Use with caution in pregnant animals Do not administer intraarterial injection, otherwise it will cause central nervous stimulation, ataxia, hyperventilation and muscle weakness.

WITHDRAWAL TIME

For cattle and Swine Meat: 28 days

STORAGE AND EXPIRED TIME

Store in a cool place and avoid sunlight.
3 years.

PACKING

50ml, 100ml.

CONCENTRATION

5.0%

IVECLO-10

COMPOSITION

Each 1ml contains

Ivermectin 10mg
Clorsulon 100mg

INDICATIONS

It is indicated for the following endo- and ectoparasites. Ivermectin and clorsulon injectable solution is an antiparasitic drug which has the broad spectrum of activity. The antiparasitic activity of ivermectin results from an increased release of neurotransmitter gamma amino butyric acid (GABA) and the flukicide activity of clorsulon results from the inhibition of enzyme systems of the parasite. Ivermectin & Clorsulon Injection has very low toxicity, has a wide safety margin and has prolonged activity. Ivermectin & Clorsulon Injection is profitable. With only one injection it's possible to control various endo- and ecto-parasites.

DOSAGE AND ADMINISTRATION


It should be administered by subcutaneous injection under the loose skin at neck region or behind the scapula. The recommended dosage levels of active substances are 0.2 mg for ivermectin and 2.0 mg for clorsulon per kg of bodyweight. The dosage level is 1 ml per 50 kg of body weight.

PRECAUTION

Do not permit water runoff from feedlots to enter lakes, streams or ponds.
Do not contaminate water by direct application or the improper disposal of drug containers.
Dispose of containers in an approved landfill or by incineration. Not for use in human's and Keep this and all drugs out of the reach of children.

WITHDRAWAL TIME

Meat: Do not use less than 28 days before slaughter for human consumption.



STORAGE AND EXPIRED TIME

Store in a dry and dark place below 30°C.
3 years.

PACKING

50ml, 100ml

CONCENTRATION

1.0%+10%

INJECTIONS

VITAB-COMP

COMPOSITION

Each 1ml contains

Vitamin B1	10mg
Vitamin B2	5mg
Vitamin B6	5mg
Vitamin B12	20mcg
Biotin	100mcg
D-Panthenol	12.5mg
Nictinamide	50mg
Choline chloride	10mg



INDICATIONS

This product is a combination of B vitamins used to treat or prevent vitamin deficiency due to poor diet, certain illnesses or during pregnancy.

DOSAGE AND ADMINISTRATION

for Intramuscular injection
Horses and cattle: a single dose of 10-20ml
Sheep: 2-3ml.

SIDE EFFECT

Vitamin B-complex includes several different components, each of which has the potential to interact with drugs.


WITHDRAWAL TIME

None.

PAKING

50ml, 100ml

TETRAOX-200



COMPOSITION

Each 1ml contains
Oxytetracycline

200mg

INDICATIONS

Treatment of diseases caused by Oxytetracycline-susceptible organisms in cattle, sheep and goats. For cattle: Bronchopneumonia and other respiratory infections, infections of the gastrointestinal tract, metritis, mastitis, septicaemia, puerperal infections, and secondary bacterial infections primarily caused by viruses, etc. For sheep and goats: Infections of respiratory, urogenital, gastrointestinal tract and hooves, mastitis, infected wounds, etc.

SIDE EFFECT AND CONTRAINDICATIONS

Administer by intramuscular injection. Cattle, sheep and goats: 0.5ml-1ml per 10kg body weight for one single dose, not more than 10ml per injection site.

ADMINISTRATION AND DOSAGE

Oxytetracycline injection is not intended for cats, dogs and horses.

It should not be given to animals in late pregnancy, animals with severe damages of liver and kidneys and to animals oversensitive to Oxytetracycline.

Sometimes a temporary swelling on the injection site occurs.

WITHDRAWAL TIME

Meat: 28 days.
Milk: 7 days.


STORAGE AND EXPIRED TIME

Store in a dry and dark place between 15°C and 25°C.
3 years.

PAKING

50ml, 100ml

TYFO-300



TREATMENT AND CONTROL OF INFECTIONS CAUSED BY ORGANISMS SENSITIVE TO FOSFOMYCIN AND TYLOSIN.

COMPOSITION

Each ml contains:
Fosfomycin 200 mg
Tylosin tartrate 100 mg

INDICATIONS

It is indicated in the treatment and control of infections caused by organisms sensitive to Fosfomycin and Tylosin. These infections include Colibacillosis, CRD Complex, infectious Coryza, Fowl Cholera and Staphylococcosis.

DOSAGE AND ADMINISTRATION

Regular dose: 1 - 0.5 g/liter/day in drinking water, for 5 - 3 days
Broiler chicken:

Age (days)	Bodyweight (g)	Approx. Water-intake liter/day/1000birds	TYFO -300 g/day/1000birds
7	170	70	35
14	400	120	80
21	730	180	120
28	1000	220	200
35	1500	270	270
42	1900	320	320

* The total amount of TYFO 300 - calculated to be dissolved in drinking water enough for 6 hours in the morning.